# CONTENTS \_\_\_\_

	We	xviii	
		xix	
		Two all-purpose study tips	XX
		Key sections for exam review (by topic)	XX
	Ack	knowledgments	xxiii
1	Fun	actions, Graphs, and Lines	1
	1.1	Functions	1
		1.1.1 Interval notation	3
		1.1.2 Finding the domain	4
		1.1.3 Finding the range using the graph	5
		1.1.4 The vertical line test	6
	1.2	Inverse Functions	7
		1.2.1 The horizontal line test	8
		1.2.2 Finding the inverse	9
		1.2.3 Restricting the domain	9
		1.2.4 Inverses of inverse functions	11
	1.3	Composition of Functions	11
	1.4	Odd and Even Functions	14
	1.5	Graphs of Linear Functions	17
	1.6	Common Functions and Graphs	19
<b>2</b>	Rev	view of Trigonometry	25
	2.1	The Basics	25
	2.2	Extending the Domain of Trig Functions	28
		2.2.1 The ASTC method	31
		2.2.2 Trig functions outside $[0, 2\pi]$	33
	2.3	The Graphs of Trig Functions	35
	2.4	Trig Identities	39

VIII • Contents

3	Inti	roduction to Limits	41
	3.1	Limits: The Basic Idea	41
	3.2	Left-Hand and Right-Hand Limits	43
	3.3	When the Limit Does Not Exist	45
	3.4	Limits at $\infty$ and $-\infty$	10
	0.4	3.4.1 Large numbers and small numbers	48
	3.5	Two Common Misconceptions about Asymptotes	50
	3.6	The Sandwich Principle	51
	3.7	Summary of Basic Types of Limits	54
4	Hoy	w to Solve Limit Problems Involving Polynomials	57
	4.1	Limits Involving Rational Functions as $x \to a$	57
	4.2	Limits Involving Square Roots as $x \to a$	61
	4.3	Limits Involving Rational Functions as $x \to \infty$	61
		4.3.1 Method and examples	64
	4.4	Limits Involving Poly-type Functions as $x \to \infty$	66
	4.5	Limits Involving Rational Functions as $x \to -\infty$	70
	4.6	Limits Involving Absolute Values	72
5	Сот	ntinuity and Differentiability	75
	5.1	Continuity	75
		5.1.1 Continuity at a point	76
		5.1.2 Continuity on an interval	77
		5.1.3 Examples of continuous functions	77
		5.1.4 The Intermediate Value Theorem	80
		5.1.5 A harder IVT example	82
		5.1.6 Maxima and minima of continuous functions	82
	5.2	Differentiability	84
		5.2.1 Average speed	84
		5.2.2 Displacement and velocity	85
		5.2.3 Instantaneous velocity	80
		5.2.4 The graphical interpretation of velocity	81
		5.2.6 The derivative function	00
		5.2.0 The derivative runction $5.2.7$ The derivative as a limiting ratio	90 91
		5.2.8 The derivative of linear functions	91
		5.2.9 Second and higher-order derivatives	94 94
		5.2.10 When the derivative does not exist	94
		5.2.11 Differentiability and continuity	96
6	Ho	w to Solve Differentiation Problems	99
	6.1	Finding Derivatives Using the Definition	99
	6.2	Finding Derivatives (the Nice Way)	102
		6.2.1 Constant multiples of functions	103

		6.2.2 Sums and differences of functions	103
		6.2.3 Products of functions via the product rule	104
		6.2.4 Quotients of functions via the quotient rule	105
		6.2.5 Composition of functions via the chain rule	107
		6.2.6 A nasty example	109
		6.2.7 Justification of the product rule and the chain rule	111
	6.3	Finding the Equation of a Tangent Line	114
	6.4	Velocity and Acceleration	114
		6.4.1 Constant negative acceleration	115
	6.5	Limits Which Are Derivatives in Disguise	117
	6.6	Derivatives of Piecewise-Defined Functions	119
	6.7	Sketching Derivative Graphs Directly	123
7	Trig	g Limits and Derivatives	127
	7.1	Limits Involving Trig Functions	127
		7.1.1 The small case	128
		7.1.2 Solving problems—the small case	129
		7.1.3 The large case	134
		7.1.4 The "other" case	137
		7.1.5 Proof of an important limit	137
	7.2	Derivatives Involving Trig Functions	141
		7.2.1 Examples of differentiating trig functions	143
		7.2.3 A curious function	145 146
		1.2.3 A curious function	140
8	Imp	blicit Differentiation and Related Rates	149
	8.1	Implicit Differentiation	149
		8.1.1 Techniques and examples	150
		8.1.2 Finding the second derivative implicitly	154
	8.2	Related Rates	156
		8.2.1 A simple example	157
		8.2.2 A slightly harder example	159
		8.2.4 A really hard example	162
		8.2.4 A rearry naru example	102
9	Exp	ponentials and Logarithms	167
	9.1	The Basics	167
		9.1.1 Review of exponentials	167
		9.1.2 Review of logarithms	100
		9.1.4 Log rules	109
	99	Definition of $e$	172
	5.4	9.2.1 A question about compound interest	173
		9.2.2 The answer to our question	173
		9.2.3 More about $e$ and logs	175
	9.3	Differentiation of Logs and Exponentials	177
		~ *	

#### $\times$ $\bullet$ Contents

		9.3.1	Examples of differentiating exponentials and logs	179
	9.4	How to	Solve Limit Problems Involving Exponentials or Logs	180
	-	9.4.1	Limits involving the definition of $e$	181
		9.4.2	Behavior of exponentials near 0	182
		9.4.3	Behavior of logarithms near 1	183
		9.4.4	Behavior of exponentials near $\infty$ or $-\infty$	184
		9.4.5	Behavior of logs near $\infty$	187
		9.4.6	Behavior of logs near 0	188
	9.5	Logari	thmic Differentiation	189
		9.5.1	The derivative of $x^a$	192
	9.6	Expon	ential Growth and Decay	193
		9.6.1	Exponential growth	194
		9.6.2	Exponential decay	195
1	9.7	Hyper	bolic Functions	198
10	Inve	erse Fu	nctions and Inverse Trig Functions	201
	10.1	The D	erivative and Inverse Functions	201
	10.1	10.1.1	Using the derivative to show that an inverse exists	201
		10.1.1 10.1.2	Derivatives and inverse functions: what can go wrong	201
		10.1.3	Finding the derivative of an inverse function	200
		10.1.4	A big example	206
	10.2	Inverse	Trig Functions	208
	10.2	10.2.1	Inverse sine	208
		10.2.2	Inverse cosine	211
		10.2.3	Inverse tangent	213
		10.2.4	Inverse secant	216
		10.2.5	Inverse cosecant and inverse cotangent	217
		10.2.6	Computing inverse trig functions	218
	10.3	Inverse	e Hyperbolic Functions	220
		10.3.1	The rest of the inverse hyperbolic functions	222
11	The	Domin	ative and Creeks	225
11	Tue	Deriv		220
	11.1	Extren	na of Functions	225
		11.1.1	Global and local extrema	225
		11.1.2 11.1.2	How to find global maxima and minima	221
	11 2	Rolle's	Theorem	220
	11.2	The M	Teon Value Theorem	- <u></u>
	11.5	11.3.1	Consequences of the Mean Value Theorem	$\frac{235}{235}$
	11 4	The Se	econd Derivative and Graphs	237
		11.4.1	More about points of inflection	238
	11 5	Classif	Tring Points Where the Derivative Vanishes	220
	11.0	11 5 1	Using the first derivative	209 240
		11.5.2	Using the second derivative	242
		· · · ·	0	

#### Contents • Xi

$12 \ \mathrm{Sk}$	etching Graphs	<b>245</b>
12	1 How to Construct a Table of Signs	245
	12.1.1 Making a table of signs for the derivative	247
	12.1.2 Making a table of signs for the second derivative	248
12	2 The Big Method	250
12	3 Examples	252
	12.3.1 An example without using derivatives	252
	12.3.2 The full method: example 1	254
	12.3.3 The full method: example 2	256
	12.3.4 The full method: example 3	259
	12.3.5 The full method: example 4	262
13 Oj	ptimization and Linearization	267
13	1 Optimization	267
	13.1.1 An easy optimization example	267
	13.1.2 Optimization problems: the general method	269
	13.1.3 An optimization example	269
	13.1.4 Another optimization example	271
	13.1.5 Using implicit differentiation in optimization	274
	13.1.6 A difficult optimization example	275
13	2 Linearization	278
	13.2.1 Linearization in general	279
	13.2.2 The differential	281
	13.2.3 Linearization summary and examples	283
	13.2.4 The error in our approximation	285
13	.3 Newton's Method	287
14 L'	Hôpital's Rule and Overview of Limits	293
14	.1 L'Hôpital's Rule	293
	14.1.1 Type <b>A</b> : $0/0$ case	294
	14.1.2 Type A: $\pm \infty / \pm \infty$ case	296
	14.1.3 Type <b>B1</b> $(\infty - \infty)$	298
	14.1.4 Type <b>B2</b> $(0 \times \pm \infty)$	299
	14.1.5 Type <b>C</b> $(1^{\pm \infty}, 0^0, \text{ or } \infty^0)$	301
	14.1.6 Summary of l'Hôpital's Rule types	302
14	2 Overview of Limits	303
15 In	troduction to Integration	307
15	1 Sigma Notation	307
10	15.1.1 A nice sum	310
	15.1.2 Telescoping series	311
15	2 Displacement and Area	21/
10	15.2.1 Three simple cases	314 314
	15.2.1 A more general journey	314
	15.2.3 Signed area	319
	15.2.4 Continuous velocity	320
	v v	

XII • Contents

	15.2.5 Two special approximations	323
16 Defi	nite Integrals	325
16.1	The Basic Idea	325
	16.1.1 Some easy examples	327
16.2	Definition of the Definite Integral	330
	16.2.1 An example of using the definition	331
16.3	Properties of Definite Integrals	334
16.4	Finding Areas	339
	16.4.1 Finding the unsigned area	339
	16.4.2 Finding the area between two curves	342
10 5	10.4.3 Finding the area between a curve and the $y$ -axis	344
16.5	Estimating Integrals	340 347
16.6	Avenues and the Mean Value Theorem for Internals	250
10.0	16.6.1 The Mean Value Theorem for integrals	351 351
16 7	A Nonintegrable Function	252
10.7	A nonintegrable Function	000
17 The	Fundamental Theorems of Calculus	355
17.1	Functions Based on Integrals of Other Functions	355
17.2	The First Fundamental Theorem	358
	17.2.1 Introduction to antiderivatives	361
17.3	The Second Fundamental Theorem	362
17.4	Indefinite Integrals	364
17.5	How to Solve Problems: The First Fundamental Theorem	366
	17.5.1 Variation 1: variable left-hand limit of integration	367
	17.5.2 Variation 2: one tricky limit of integration	367
	17.5.3 Variation 3: two tricky limits of integration	369 270
17.6	17.5.4 Variation 4: mint is a derivative in disguise	970 971
17.0	How to Solve Problems: The Second Fundamental Theorem	371
	17.6.2 Finding definite integrals	374
	17.6.3 Unsigned areas and absolute values	376
17.7	A Technical Point	380
17.8	Proof of the First Fundamental Theorem	381
18 Tech	niques of Integration. Part One	383
18.1	Substitution	383
10.1	18.1.1 Substitution and definite integrals	386
	18.1.2 How to decide what to substitute	389
	18.1.3 Theoretical justification of the substitution method	392
18.2	Integration by Parts	393
	18.2.1 Some variations	394
18.3	Partial Fractions	397

#### Contents • XIII

18.3.1 The algebra of partial fractions	398
18.3.2 Integrating the pieces	401
18.3.3 The method and a big example	404
19 Techniques of Integration, Part Two	409
19.1 Integrals Involving Trig Identities	409
19.2 Integrals Involving Powers of Trig Functions	413
19.2.1 Powers of $\sin$ and/or $\cos$	413
19.2.2 Powers of tan	415
19.2.3 Powers of sec	416
19.2.4 Powers of cot	418
19.2.5 Fowers of CSC 19.2.6 Reduction formulas	410
10.3 Integrals Involving Trig Substitutions	491
19.3 Integrals involving Trig Substitutions 10.3.1 Type 1: $\sqrt{a^2 - x^2}$	421
19.3.1 Type 1. $\sqrt{a^2 - x^2}$	421
19.3.2 Type 2. $\sqrt{x + u}$ 10.3.3 Type 2. $\sqrt{x^2 - a^2}$	420
19.3.4 Completing the square and trig substitutions	424
19.3.5 Summary of trig substitutions	426
19.3.6 Technicalities of square roots and trig substitutions	427
19.4 Overview of Techniques of Integration	429
20 Improper Integrals: Basic Concepts	431
20.1 Convergence and Divergence	431
20.1.1 Some examples of improper integrals	433
20.1.2 Other blow-up points	435
20.2 Integrals over Unbounded Regions	437
20.3 The Comparison Test (Theory)	439
20.4 The Limit Comparison Test (Theory)	441
20.4.1 Functions asymptotic to each other	441
20.4.2 The statement of the test	443
20.5 The $p$ -test (Theory)	444
20.6 The Absolute Convergence Test	447
21 Improper Integrals: How to Solve Problems	451
21.1 How to Get Started	451
21.1.1 Splitting up the integral	452
21.1.2 How to deal with negative function values	453
21.2 Summary of Integral Tests	454
21.3 Behavior of Common Functions near $\infty$ and $-\infty$	456
21.3.1 Polynomials and poly-type functions near $\infty$ and $-\infty$	456
21.3.2 Trig functions near $\infty$ and $-\infty$	459
21.3.3 Exponentials near $\infty$ and $-\infty$	461
21.3.4 Logarithms near $\infty$	465
21.4 Behavior of Common Functions near 0	469

#### $XiV \bullet Contents$

21.4.1 Polynomials and poly-type functions near $0$	469
21.4.2 Trig functions near 0	470
21.4.3 Exponentials near 0	472
21.4.4 Logarithms near 0	473
21.4.5 The behavior of more general functions near U	474
21.5 How to Deal with Problem Spots Not at 0 or $\infty$	475
22 Sequences and Series: Basic Concepts	477
22.1 Convergence and Divergence of Sequences	477
22.1.1 The connection between sequences and functions	478
22.1.2 Two important sequences	480
22.2 Convergence and Divergence of Series	481
22.2.1 Geometric series (theory)	484
22.3 The <i>n</i> th Term Test (Theory)	486
22.4 Properties of Both Infinite Series and Improper Integrals	487
22.4.1 The comparison test (theory)	487
22.4.2 The limit comparison test (theory)	488
22.4.3 The $p$ -test (theory)	489
22.4.4 The absolute convergence test	490
22.5 New Tests for Series	491
22.5.1 The ratio test (theory)	492
22.5.2 The root test (theory)	493
22.5.3 The integral test (theory)	494
22.5.4 The alternating series test (theory)	497
23 How to Solve Series Problems	501
23.1 How to Evaluate Geometric Series	502
23.2 How to Use the $n$ th Term Test	503
23.3 How to Use the Ratio Test	504
23.4 How to Use the Root Test	508
23.5 How to Use the Integral Test	509
23.6 Comparison Test, Limit Comparison Test, and $p$ -test	510
23.7 How to Deal with Series with Negative Terms	515
24 Taylor Polynomials, Taylor Series, and Power Series	519
24.1 Approximations and Taylor Polynomials	519
24.1.1 Linearization revisited	520
24.1.2 Quadratic approximations	521
24.1.3 Higher-degree approximations	522
24.1.4 Taylor's Theorem	523
24.2 Power Series and Taylor Series	526
24.2.1 Power series in general	527
24.2.2 Taylor series and Maclaurin series	529
24.2.3 Convergence of Taylor series	530

#### Contents $\bullet \times \vee$

25 How to Solve Estimation Problems	535
25.1 Summary of Taylor Polynomials and Series	535
25.2 Finding Taylor Polynomials and Series	537
25.3 Estimation Problems Using the Error Term	540
25.3.1 First example	541
25.3.2 Second example	543
25.3.3 Third example	544
25.3.4 Fourth example	546
25.3.5 Fifth example	547
25.3.6 General techniques for estimating the error term	548
25.4 Another Technique for Estimating the Error	548
26 Taylor and Power Series: How to Solve Problems	551
26.1 Convergence of Power Series	551
26.1.1 Radius of convergence	551
26.1.2 How to find the radius and region of convergence	554
26.2 Getting New Taylor Series from Old Ones	558
26.2.1 Substitution and Taylor series	560
26.2.2 Differentiating Taylor series	562
26.2.3 Integrating Taylor series	563
26.2.4 Adding and subtracting Taylor series	565
26.2.5 Multiplying Taylor series	500 567
26.2.0 Dividing Taylor Series	568
26.4 Using Maclaurin Series to Find Limits	508 570
2011 OSing Madaalin Solids to Tina Linnis	010
27 Parametric Equations and Polar Coordinates	575
27.1 Parametric Equations	575
27.1.1 Derivatives of parametric equations	578
27.2 Polar Coordinates	581
27.2.1 Converting to and from polar coordinates	582
27.2.2 Sketching curves in polar coordinates	585
27.2.3 Finding tangents to polar curves	590
27.2.4 Finding areas enclosed by polar curves	591
28 Complex Numbers	595
28.1 The Basics	595
28.1.1 Complex exponentials	598
28.2 The Complex Plane	599
28.2.1 Converting to and from polar form	601
28.3 Taking Large Powers of Complex Numbers	603
28.4 Solving $z^n = w$	604
28.4.1 Some variations	608
28.5 Solving $e^z = w$	610
28.6 Some Trigonometric Series	612

XVI • Contents

28.7	Euler's Identity and Power Series	615
29 Vol	umes, Arc Lengths, and Surface Areas	617
29.1	Volumes of Solids of Revolution	617
	29.1.1 The disc method	619
	29.1.2 The shell method	620
	29.1.3 Summary and variations	622
	29.1.4 Variation 1: regions between a curve and the $y$ -axis	623
	29.1.5 Variation 2: regions between two curves	625
	29.1.6 Variation 3: axes parallel to the coordinate axes	628
29.2	Volumes of General Solids	631
29.3	Arc Lengths	637
	29.3.1 Parametrization and speed	639
29.4	Surface Areas of Solids of Revolution	640
30 Diff	Cerential Equations	645
30.1	Introduction to Differential Equations	645
30.2	Separable First-order Differential Equations	646
30.3	First-order Linear Equations	648
	30.3.1 Why the integrating factor works	652
30.4	Constant-coefficient Differential Equations	653
	30.4.1 Solving first-order homogeneous equations	654
	30.4.2 Solving second-order homogeneous equations	654
	30.4.3 Why the characteristic quadratic method works	655
	30.4.4 Nonhomogeneous equations and particular solutions	656
	30.4.5 Finding a particular solution	658
	30.4.6 Examples of finding particular solutions	660
	30.4.7 Resolving conflicts between $y_P$ and $y_H$	662
	30.4.8 Initial value problems (constant-coefficient linear)	663
30.5	Modeling Using Differential Equations	665
Appen	dix A Limits and Proofs	669
A.1	Formal Definition of a Limit	669
	A.1.1 A little game	670
	A.1.2 The actual definition	672
	A.1.3 Examples of using the definition	672
A.2	Making New Limits from Old Ones	674
	A.2.1 Sums and differences of limits—proofs	674
	A.2.2 Products of limits—proof	675
	A.2.3 Quotients of limits—proof	676
	A.2.4 The sandwich principle—proof	678
A.3	Other Varieties of Limits	678
	A.3.1 Infinite limits	679
	A.3.2 Left-hand and right-hand limits	680
	A.3.3 Limits at $\infty$ and $-\infty$	680

#### Contents • XVII

	A.3.4 Two examples involving trig	682
A.4	Continuity and Limits	684
	A.4.1 Composition of continuous functions	684
	A.4.2 Proof of the Intermediate Value Theorem	686
	A.4.3 Proof of the Max-Min Theorem	687
A.5	Exponentials and Logarithms Revisited	689
A.6	Differentiation and Limits	691
	A.6.1 Constant multiples of functions	691
	A.6.2 Sums and differences of functions	691
	A.6.3 Proof of the product rule	692
	A.6.4 Proof of the quotient rule	693
	A.6.5 Proof of the chain rule	693
	A.6.6 Proof of the Extreme Value Theorem	694
	A.6.7 Proof of Rolle's Theorem	695
	A.6.8 Proof of the Mean Value Theorem	695
	A.6.9 The error in linearization	696
	A.6.10 Derivatives of piecewise-defined functions	697
	A.6.11 Proof of l'Hôpital's Rule	698
A.7	Proof of the Taylor Approximation Theorem	700
Appen	dix B Estimating Integrals	703
B.1	Estimating Integrals Using Strips	703
	B.1.1 Evenly spaced partitions	705
B.2	The Trapezoidal Rule	706
B.3	Simpson's Rule	709
	B.3.1 Proof of Simpson's rule	710
B.4	The Error in Our Approximations	711
	B.4.1 Examples of estimating the error	712
	B.4.2 Proof of an error term inequality	714
$\mathbf{List}$	of Symbols	717
Inde	ex	719

# CHAPTER 1

# Functions, Graphs, and Lines

Trying to do calculus without using functions would be one of the most pointless things you could do. If calculus had an ingredients list, functions would be first on it, and by some margin too. So, the first two chapters of this book are designed to jog your memory about the main features of functions. This chapter contains a review of the following topics:

- functions: their domain, codomain, and range, and the vertical line test;
- inverse functions and the horizontal line test;
- composition of functions;
- odd and even functions;
- graphs of linear functions and polynomials in general, as well as a brief survey of graphs of rational functions, exponentials, and logarithms; and
- how to deal with absolute values.

Trigonometric functions, or trig functions for short, are dealt with in the next chapter. So, let's kick off with a review of what a function actually is.

## 1.1 Functions

A *function* is a rule for transforming an object into another object. The object you start with is called the *input*, and comes from some set called the *domain*. What you get back is called the *output*; it comes from some set called the *codomain*.

Here are some examples of functions:

• Suppose you write  $f(x) = x^2$ . You have just defined a function f which transforms any number into its square. Since you didn't say what the domain or codomain are, it's assumed that they are both  $\mathbb{R}$ , the set of all real numbers. So you can square any real number, and get a real number back. For example, f transforms 2 into 4; it transforms -1/2 into 1/4; and it transforms 1 into 1. This last one isn't much of a change at all, but that's no problem: the transformed object doesn't have to be different from the original one. When you write f(2) = 4, what you really mean

2 • Functions, Graphs, and Lines

is that f transforms 2 into 4. By the way, f is the transformation rule, while f(x) is the result of applying the transformation rule to the variable x. So it's technically not correct to say "f(x) is a function"; it should be "f is a function."

- Now, let  $g(x) = x^2$  with domain consisting only of numbers greater than or equal to 0. (Such numbers are called *nonnegative*.) This seems like the same function as f, but it's not: the domains are different. For example, f(-1/2) = 1/4, but g(-1/2) isn't defined. The function g just chokes on anything not in the domain, refusing even to touch it. Since g and f have the same rule, but the domain of g is smaller than the domain of f, we say that g is formed by *restricting the domain* of f.
- Still letting  $f(x) = x^2$ , what do you make of f(horse)? Obviously this is undefined, since you can't square a horse. On the other hand, let's set

$$h(x) =$$
 number of legs x has,

where the domain of h is the set of all animals. So h(horse) = 4, while h(ant) = 6 and h(salmon) = 0. The codomain could be the set of all nonnegative integers, since animals don't have negative or fractional numbers of legs. By the way, what is h(2)? This isn't defined, of course, since 2 isn't in the domain. How many legs does a "2" have, after all? The question doesn't really make sense. You might also think that h(chair) = 4, since most chairs have four legs, but that doesn't work either, since a chair isn't an animal, and so "chair" is not in the domain of h. That is, h(chair) is undefined.

• Suppose you have a dog called Junkster. Unfortunately, poor Junkster has indigestion. He eats something, then chews on it for a while and tries to digest it, fails, and hurls. Junkster has transformed the food into ... something else altogether. We could let

j(x) =color of barf when Junkster eats x,

where the domain of j is the set of foods that Junkster will eat. The codomain is the set of all colors. For this to work, we have to be confident that whenever Junkster eats a taco, his barf is always the same color (say, red). If it's sometimes red and sometimes green, that's no good: **a** function must assign a unique output for each valid input.

Now we have to look at the concept of the *range* of a function. The range is the set of all outputs that could possibly occur. You can think of the function working on transforming everything in the domain, one object at a time; the collection of transformed objects is the range. You might get duplicates, but that's OK.

So why isn't the range the same thing as the codomain? Well, the range is actually a subset of the codomain. The codomain is a set of **possible** outputs, while the range is the set of **actual** outputs. Here are the ranges of the functions we looked at above:

#### Section 1.1.1: Interval notation • 3

- If  $f(x) = x^2$  with domain  $\mathbb{R}$  and codomain  $\mathbb{R}$ , the range is the set of nonnegative numbers. After all, when you square a number, the result cannot be negative. How do you know the range is **all** the nonnegative numbers? Well, if you square every number, you definitely cover all nonnegative numbers. For example, you get 2 by squaring  $\sqrt{2}$  (or  $-\sqrt{2}$ ).
- If  $g(x) = x^2$ , where the domain of g is only the nonnegative numbers but the codomain is still all of  $\mathbb{R}$ , the range will again be the set of nonnegative numbers. When you square every nonnegative number, you still cover all the nonnegative numbers.
- If h(x) is the number of legs the animal x has, then the range is all the possible numbers of legs that **any** animal can have. I can think of animals that have 0, 2, 4, 6, and 8 legs, as well as some creepy-crawlies with more legs. If you include individual animals which have lost one or more legs, you can also include 1, 3, 5, and 7 in the mix, as well as other possibilities. In any case, the range of this function isn't so clear-cut; you probably have to be a biologist to know the real answer.
- Finally, if j(x) is the color of Junkster's barf when he eats x, then the range consists of all possible barf-colors. I dread to think what these are, but probably bright blue isn't among them.

#### 1.1.1 Interval notation

In the rest of this book, our functions will always have codomain  $\mathbb{R}$ , and the domain will always be as much of  $\mathbb{R}$  as possible (unless stated otherwise). So we'll often be dealing with subsets of the real line, especially connected intervals such as  $\{x : 2 \le x < 5\}$ . It's a bit of a pain to write out the full set notation like this, but it sure beats having to say "all the numbers between 2 and 5, including 2 but not 5." We can do even better using interval notation.

We'll write [a, b] to mean the set of all numbers between a and b, including a and b themselves. So [a, b] means the set of all x such that  $a \le x \le b$ . For example, [2, 5] is the set of all real numbers between 2 and 5, including 2 and 5. (It's not just the set consisting of 2, 3, 4, and 5: don't forget that there are loads of fractions and irrational numbers between 2 and 5, such as 5/2,  $\sqrt{7}$ , and  $\pi$ .) An interval such as [a, b] is called *closed*.

If you don't want the endpoints, change the square brackets to parentheses. In particular, (a, b) is the set of all numbers between a and b, not including a or b. So if x is in the interval (a, b), we know that a < x < b. The set (2, 5) includes all real numbers between 2 and 5, but not 2 or 5. An interval of the form (a, b) is called *open*.

You can mix and match: [a, b) consists of all numbers between a and b, including a but not b. And (a, b] includes b but not a. These intervals are closed at one end and open at the other. Sometimes such intervals are called *half-open*. An example is the set  $\{x : 2 \le x < 5\}$  from above, which can also be written as [2, 5).

There's also the useful notation  $(a, \infty)$  for all the numbers greater than a not including a;  $[a, \infty)$  is the same thing but with a included. There are three other possibilities which involve  $-\infty$ ; all in all, the situation looks like this:

4 • Functions, Graphs, and Lines



#### 1.1.2 Finding the domain

Sometimes the definition of a function will include the domain. (This was the case, for example, with our function g from Section 1.1 above.) Most of the time, however, the domain is not provided. The basic convention is that the domain consists of as much of the set of real numbers as possible. For example, if  $k(x) = \sqrt{x}$ , the domain can't be all of  $\mathbb{R}$ , since you can't take the square root of a negative number. The domain must be  $[0, \infty)$ , which is just the set of all numbers greater than or equal to 0.

OK, so square roots of negative numbers are bad. What else can cause a screw-up? Here's a list of the three most common possibilities:

- 1. The denominator of a fraction can't be zero.
- 2. You can't take the square root (or fourth root, sixth root, and so on) of a negative number.
- 3. You can't take the logarithm of a negative number or of 0. (Remember logs? If not, see Chapter 9!)

You might recall that  $\tan(90^\circ)$  is also a problem, but this is really a special case of the first item above. You see,

$$\tan(90^{\circ}) = \frac{\sin(90^{\circ})}{\cos(90^{\circ})} = \frac{1}{0},$$



so the reason  $\tan(90^\circ)$  is undefined is really that a hidden denominator is zero. Here's another example: if we try to define

$$f(x) = \frac{\log_{10}(x+8)\sqrt{26-2x}}{(x-2)(x+19)},$$

then what is the domain of f? Well, for f(x) to make sense, here's what needs to happen:

• We need to take the square root of (26-2x), so this quantity had better be nonnegative. That is,  $26-2x \ge 0$ . This can be rewritten as  $x \le 13$ .

Section 1.1.3: Finding the range using the graph  $\bullet$  5

- We also need to take the logarithm of (x + 8), so this quantity needs to be positive. (Notice the difference between logs and square roots: you can take the square root of 0, but you can't take the log of 0.) Anyway, we need x + 8 > 0, so x > -8. So far, we know that  $-8 < x \le 13$ , so the domain is at most (-8, 13].
- The denominator can't be 0; this means that  $(x-2) \neq 0$  and  $(x+19) \neq 0$ . In other words,  $x \neq 2$  and  $x \neq -19$ . This last one isn't a problem, since we already know that x lies in (-8, 13], so x can't possibly be -19. We do have to exclude 2, though.

So we have found that the domain is the set (-8, 13] except for the number 2. This set could be written as  $(-8, 13] \setminus \{2\}$ . Here the backslash means "not including."

#### 1.1.3 Finding the range using the graph

Let's define a new function F by specifying that its domain is [-2, 1] and that  $F(x) = x^2$  on this domain. (Remember, the codomain of any function we look at will always be the set of all real numbers.) Is F the same function as f, where  $f(x) = x^2$  for all real numbers x? The answer is no, since the two functions have different domains (even though they have the same rule). As in the case of the function g from Section 1.1 above, the function F is formed by restricting the domain of f.

Now, what is the range of F? Well, what happens if you square every number between -2 and 1 inclusive? You should be able to work this out directly, but this is a good opportunity to see how to use a graph to find the range of a function. The idea is to sketch the graph of the function, then imagine two rows of lights shining from the far left and far right of the graph horizontally toward the y-axis. The curve will cast two shadows, one on the left side and one on the right side of the y-axis. The range is the union of both shadows: that is, if any point on the y-axis lies in either the left-hand or the right-hand shadow, it is in the range of the function. Let's see how this works with our function F:



6 • Functions, Graphs, and Lines

The left-hand shadow covers all the points on the *y*-axis between 0 and 4 inclusive, which is [0, 4]; on the other hand, the right-hand shadow covers the points between 0 and 1 inclusive, which is [0, 1]. The right-hand shadow doesn't contribute anything extra: the total coverage is still [0, 4]. This is the range of F.

1.1.4 The vertical line test

In the last section, we used the graph of a function to find its range. The graph of a function is very important: it really shows you what the function "looks like." We'll be looking at techniques for sketching graphs in Chapter 12, but for now I'd like to remind you about the vertical line test.

You can draw any figure you like on a coordinate plane, but the result may not be the graph of a function. So what's special about the graph of a function? What is the graph of a function f, anyway? Well, it's the collection of all points with coordinates (x, f(x)), where x is in the domain of f. Here's another way of looking at this: start with some number x. If x is in the domain, you plot the point (x, f(x)), which of course is at a height of f(x)units above the point x on the x-axis. If x isn't in the domain, you don't plot anything. Now repeat for every real number x to build up the graph.

Here's the key idea: you can't have two points with the same x-coordinate. In other words, no two points on the graph can lie on the same vertical line. Otherwise, how would you know which of the two or more heights above the point x on the x-axis corresponds to the value of f(x)? So, this leads us to the vertical line test: if you have some graph and you want to know whether it's the graph of a function, see whether any vertical line intersects the graph more than once. If so, it's not the graph of a function; but if no vertical line intersects the graph more than once, you are indeed dealing with the graph of a function. For example, the circle of radius 3 units centered at the origin has a graph like this:



Such a commonplace object should be a function, right? No, check the vertical lines that are shown in the diagram. Sure, to the left of -3 or to the right of 3, there's no problem—the vertical lines don't even hit the graph, which is fine. Even at -3 or 3, the vertical lines only intersect the curve in one point each, which is also fine. The problem is when x is in the interval (-3, 3). For

#### Section 1.2: Inverse Functions • 7

any of these values of x, the vertical line through (x, 0) intersects the circle twice, which screws up the circle's potential function-status. You just don't know whether f(x) is the top point or the bottom point.

The best way to salvage the situation is to chop the circle in half horizontally and choose only the top or the bottom half. The equation for the whole circle is  $x^2 + y^2 = 9$ , whereas the equation for the top semicircle is  $y = \sqrt{9 - x^2}$ . The bottom semicircle has equation  $y = -\sqrt{9 - x^2}$ . These last two are functions, both with domain [-3,3]. If you felt like chopping in a different way, you wouldn't actually have to take semicircles—you could chop and change between the upper and lower semicircles, as long as you don't violate the vertical line test. For example, here's the graph of a function which also has domain [-3,3]:



The vertical line test checks out, so this is indeed the graph of a function.

### 1.2 Inverse Functions

Let's say you have a function f. You present it with an input x; provided that x is in the domain of f, you get back an output, which we call f(x). Now we try to do things all backward and ask this question: if you pick a number y, what input can you give to f in order to get back y as your output?

Here's how to state the problem in math-speak: given a number y, what x in the domain of f satisfies f(x) = y? The first thing to notice is that y has to be in the range of f. Otherwise, by definition there are no values of x such that f(x) = y. There would be nothing in the domain that f would transform into y, since the range is **all** the possible outputs.

On the other hand, if y is in the range, there might be many values that work. For example, if  $f(x) = x^2$  (with domain  $\mathbb{R}$ ), and we ask what value of x transforms into 64, there are obviously two values of x: 8 and -8. On the other hand, if  $g(x) = x^3$ , and we ask the same question, there's only one value of x, which is 4. The same would be true for any number we give to g to transform, because any number has only one (real) cube root.

So, here's the situation: we're given a function f, and we pick y in the range of f. Ideally, there will be exactly one value of x which satisfies f(x) = y. If this is true for every value of y in the range, then we can define a new

8 • Functions, Graphs, and Lines

function which reverses the transformation. Starting with the output y, the new function finds the one and only input x which leads to the output. The new function is called the *inverse function of* f, and is written as  $f^{-1}$ . Here's a summary of the situation in mathematical language:

- 1. Start with a function f such that for any y in the range of f, there is exactly one number x such that f(x) = y. That is, different inputs give different outputs. Now we will define the inverse function  $f^{-1}$ .
- 2. The domain of  $f^{-1}$  is the same as the range of f.
- 3. The range of  $f^{-1}$  is the same as the domain of f.
- 4. The value of  $f^{-1}(y)$  is the number x such that f(x) = y. So,

if 
$$f(x) = y$$
, then  $f^{-1}(y) = x$ .

The transformation  $f^{-1}$  acts like an undo button for f: if you start with x and transform it into y using the function f, then you can undo the effect of the transformation by using the inverse function  $f^{-1}$  on y to get x back.

This raises some questions: how do you see if there's only one value of x that satisfies the equation f(x) = y? If so, how do you find the inverse, and what does its graph look like? If not, how do you salvage the situation? We'll answer these questions in the next three sections.

#### 1.2.1 The horizontal line test

For the first question—how to see that there's only one value of x that works for any y in the range—perhaps the best way is to look at the graph of your function. We want to pick y in the range of f and hopefully only have one value of x such that f(x) = y. What this means is that the horizontal line through the point (0, y) should intersect the graph exactly once, at some point (x, y). That x is the one we want. If the horizontal line intersects the curve more than once, there would be multiple potential inverses x, which is bad. In that case, the only way to get an inverse function is to restrict the domain; we'll come back to this very shortly. What if the horizontal line doesn't intersect the curve at all? Then y isn't in the range after all, which is OK.

So, we have just described the *horizontal line test*: if every horizontal line intersects the graph of a function at most once, the function has an inverse. If even one horizontal line intersects the graph more than once, there isn't an inverse function. For example, look at the graphs of  $f(x) = x^3$  and  $g(x) = x^2$ :



#### Section 1.2.2: Finding the inverse $\bullet$ 9

No horizontal line hits y = f(x) more than once, so f has an inverse. On the other hand, some of the horizontal lines hit the curve y = g(x) twice, so g has no inverse. Here's the problem: if you want to solve  $y = x^2$  for x, where y is positive, then there are two solutions,  $x = \sqrt{y}$  and  $x = -\sqrt{y}$ . You don't know which one to take!

#### 1.2.2 Finding the inverse

Now let's move on to the second of our questions: how do you find the inverse of a function f? Well, you write down y = f(x) and try to solve for x. In our example of  $f(x) = x^3$ , we have  $y = x^3$ , so  $x = \sqrt[3]{y}$ . This means that  $f^{-1}(y) = \sqrt[3]{y}$ . If the variable y here offends you, by all means switch it to x: you can write  $f^{-1}(x) = \sqrt[3]{x}$  if you prefer. Of course, solving for x is not always easy and in fact is often impossible. On the other hand, if you know what the graph of your function looks like, the graph of the inverse function is easy to find. The idea is to draw the line y = x on the graph, then pretend that this line is a two-sided mirror. The inverse function is the reflection of the original function in this mirror. When  $f(x) = x^3$ , here's what  $f^{-1}$  looks like:



The original function f is reflected in the mirror y = x to get the inverse function. Note that the domain and range of both f and  $f^{-1}$  are the whole real line.

### 1.2.3 Restricting the domain

Finally, we'll address our third question: if the horizontal line test fails and there's no inverse, what can be done? Our problem is that there are multiple values of x that give the same y. The only way to get around the problem is to throw away all but one of these values of x. That is, we have to decide which one of our values of x we want to keep, and throw the rest away. As we saw in Section 1.1 above, this is called *restricting the domain* of our function. Effectively, we ghost out part of the curve so that what's left no longer fails the horizontal line test. For example, if  $g(x) = x^2$ , we can ghost out the left half of the graph like this:

10 • Functions, Graphs, and Lines



The new (unghosted) curve has the reduced domain  $[0, \infty)$  and satisfies the horizontal line test, so there is an inverse function. More precisely, the function h, which has domain  $[0, \infty)$  and is defined by  $h(x) = x^2$  on this domain, has an inverse. Let's play the reflection game to see what it looks like:



To find the equation of the inverse, we have to solve for x in the equation  $y = x^2$ . Clearly the solution is  $x = \sqrt{y}$  or  $x = -\sqrt{y}$ , but which one do we need? We know that the range of the inverse function is the same as the domain of the original function, which we have restricted to be  $[0, \infty)$ . So we need a nonnegative number as our answer, and that has to be  $x = \sqrt{y}$ . That is,  $h^{-1}(y) = \sqrt{y}$ . Of course, we could have ghosted out the right half of the original graph to restrict the domain to  $(-\infty, 0]$ . In that case, we'd get a function j which has domain  $(-\infty, 0]$  and again satisfies  $j(x) = x^2$ , but only on this domain. This function also has an inverse, but the inverse is now the negative square root:  $j^{-1}(y) = -\sqrt{y}$ .

By the way, if you take the original function g given by  $g(x) = x^2$  with domain  $(-\infty, \infty)$ , which fails the horizontal line test, and try to reflect it in the mirror y = x, you get the following picture:



#### Section 1.2.4: Inverses of inverse functions $\bullet$

Notice that the graph fails the vertical line test, so it's not the graph of a function. This illustrates the connection between the vertical and horizontal line tests—when horizontal lines are reflected in the mirror y = x, they become vertical lines.

#### 1.2.4 Inverses of inverse functions

One more thing about inverse functions: if f has an inverse, it's true that  $f^{-1}(f(x)) = x$  for all x in the domain of f, and also that  $f(f^{-1}(y)) = y$  for all y in the range of f. (Remember, the range of f is the same as the domain of  $f^{-1}$ , so you can indeed take  $f^{-1}(y)$  for y in the range of f without causing any screwups.)



For example, if  $f(x) = x^3$ , then f has an inverse given by  $f^{-1}(x) = \sqrt[3]{x}$ , and so  $f^{-1}(f(x)) = \sqrt[3]{x^3} = x$  for any x. Remember, the inverse function is like an undo button. We use x as an input to f, and then give the output to  $f^{-1}$ ; this undoes the transformation and gives us back x, the original number. Similarly,  $f(f^{-1}(y)) = (\sqrt[3]{y})^3 = y$ . So  $f^{-1}$  is the inverse function of f, and f is the inverse function of  $f^{-1}$ . In other words, the inverse of the inverse is the original function.

Now, you have to be careful in the case where you restrict the domain. Let  $g(x) = x^2$ ; we've seen that you need to restrict the domain to get an inverse. Let's say we restrict the domain to  $[0, \infty)$  and carelessly continue to refer to the function as g instead of h, as in the previous section. We would then say that  $g^{-1}(x) = \sqrt{x}$ . If you calculate  $g(g^{-1}(x))$ , you find that this is  $(\sqrt{x})^2$ , which equals x, provided that  $x \ge 0$ . (Otherwise you can't take the square root in the first place.)

On the other hand, if you work out  $g^{-1}(g(x))$ , you get  $\sqrt{x^2}$ , which is not always the same thing as x. For example, if x = -2, then  $x^2 = 4$  and so  $\sqrt{x^2} = \sqrt{4} = 2$ . So it's not true in general that  $g^{-1}(g(x)) = x$ . The problem is that -2 isn't in the restricted-domain version of g. Technically, you can't even compute g(-2), since -2 is no longer in the domain of g. We really should be working with h, not g, so that we remember to be more careful. Nevertheless, in practice, mathematicians will often restrict the domain without changing letters! So it will be useful to summarize the situation as follows:



If the domain of a function f can be restricted so that f has an inverse  $f^{-1}$ , then

- $f(f^{-1}(y)) = y$  for all y in the range of f; but
- $f^{-1}(f(x))$  may not equal x; in fact,  $f^{-1}(f(x)) = x$  only when x is in the restricted domain.

We'll be revisiting these important points in the context of inverse trig functions in Section 10.2.6 of Chapter 10.

## 1.3 Composition of Functions

Let's say we have a function g given by  $g(x) = x^2$ . You can replace x by anything you like, as long as it makes sense. For example, you can write

12 • Functions, Graphs, and Lines

 $g(y) = y^2$ , or  $g(x+5) = (x+5)^2$ . This last example shows that you need to be very careful with parentheses. It would be wrong to write  $g(x+5) = x+5^2$ , since this is just x + 25, which is not the same thing as  $(x+5)^2$ . If in doubt, use parentheses. That is, if you need to write out f(something), replace every instance of x by (something), making sure to include the parentheses. Just about the only time you don't need to use parentheses is when the function is an exponential function—for example, if  $h(x) = 3^x$ , then you can just write  $h(x^2+6) = 3^{x^2+6}$ . You don't need parentheses since you're already writing the  $x^2 + 6$  as a superscript.

Now consider the function f defined by  $f(x) = \cos(x^2)$ . If I give you a number x, how do you compute f(x)? Well, first you square it, then you take the cosine of the result. Since we can decompose the action of f(x) into these two separate actions which are performed one after the other, we might as well describe those actions as functions themselves. So, let  $g(x) = x^2$  and  $h(x) = \cos(x)$ . To simulate what f does when you use x as an input, you could first give x to q to square it, and then instead of taking the result back you could ask q to give its result to h instead. Then h spits out a number, which is the final answer. The answer will, of course, be the cosine of what came out of q, which was the square of the original x. This behavior exactly mimics f, so we can write f(x) = h(g(x)). Another way of expressing this is to write  $f = h \circ q$ ; here the circle means "composed with." That is, f is h composed with g, or in other words, f is the *composition* of h and g. What's tricky is that you write h before q (reading from left to right as usual!) but you apply q first. I agree that it's confusing, but what can I say—you just have to deal with it.



It's useful to practice composing two or more functions together. For example, if  $g(x) = 2^x$ ,  $h(x) = 5x^4$ , and j(x) = 2x - 1, what is a formula for the function  $f = g \circ h \circ j$ ? Well, just replace one thing at a time, starting with j, then h, then g. So:

$$f(x) = g(h(j(x))) = g(h(2x-1)) = g(5(2x-1)^4) = 2^{5(2x-1)^4}.$$



You should also practice reversing the process. For example, suppose you start off with

$$f(x) = \frac{1}{\tan(5\log_2(x+3))}$$

How would you decompose f into simpler functions? Zoom in to where you see the quantity x. The first thing you do is add 3, so let g(x) = x + 3. Then you have to take the base 2 logarithm of the resulting quantity, so set  $h(x) = \log_2(x)$ . Next, multiply by 5, so set j(x) = 5x. Then take the tangent, so put  $k(x) = \tan(x)$ . Finally, take reciprocals, so let m(x) = 1/x. With all these definitions, you should check that

$$f(x) = m(k(j(h(g(x))))).$$

Using the composition notation, you can write

$$f = m \circ k \circ j \circ h \circ g.$$

#### Section 1.3: Composition of Functions • 13

This isn't the only way to break down f. For example, we could have combined h and j into another function n, where  $n(x) = 5 \log_2(x)$ . Then you should check that  $n = j \circ h$ , and

$$f = m \circ k \circ n \circ g.$$

Perhaps the original decomposition (involving j and h) is better because it breaks down f into more elementary steps, but the second one (involving n) isn't wrong. After all,  $n(x) = 5 \log_2(x)$  is still a pretty simple function of x.

Beware: composition of functions isn't the same thing as multiplying them together. For example, if  $f(x) = x^2 \sin(x)$ , then f is not the composition of two functions. To calculate f(x) for any given x, you actually have to find both  $x^2$  and  $\sin(x)$  (it doesn't matter which one you find first, unlike with composition) and then multiply these two things together. If  $g(x) = x^2$  and  $h(x) = \sin(x)$ , then we'd write f(x) = g(x)h(x), or f = gh. Compare this to the composition of the two functions,  $j = g \circ h$ , which is given by

$$j(x) = g(h(x)) = g(\sin(x)) = (\sin(x))^2$$

or simply  $j(x) = \sin^2(x)$ . The function j is a completely different function from the product  $x^2 \sin(x)$ . It's also different from the function  $k = h \circ g$ , which is also a composition of g and h but in the other order:

$$k(x) = h(g(x)) = h(x^2) = \sin(x^2).$$

This is yet another completely different function. The moral of the story is that products and compositions are not the same thing, and furthermore, the order of the functions matters when you compose them, but not when you multiply them together.

One simple but important example of composition of functions occurs when you compose some function f with g(x) = x - a, where a is some constant number. You end up with a new function h given by h(x) = f(x-a). A useful point to note is that the graph of y = h(x) is the same as the graph of y = f(x), except that it's shifted over a units to the right. If a is negative, then the shift is to the left. (The way to think of this, for example, is that a shift of -3 units to the right is the same as a shift of 3 units to the left.) So, how would you sketch the graph of  $y = (x - 1)^2$ ? This is the same as  $y = x^2$ , but with x replaced by x - 1. So the graph of  $y = x^2$  needs to be shifted to the right by 1 unit, and looks like this:





#### 14 • Functions, Graphs, and Lines

Similarly, the graph of  $y = (x+2)^2$  is the graph of  $y = x^2$  shifted to the left by 2 units, since you can interpret (x+2) as (x-(-2)).

# 1.4 Odd and Even Functions

Some functions have some symmetry properties that make them easier to deal with. Consider the function f given by  $f(x) = x^2$ . Pick any positive number you like (I'll choose 3) and hit it with f (I get 9). Now take the negative of that number, -3 in my case, and hit that with f (I get 9 again). You should get the same answer both times, as I did, regardless of which number you chose. You can express this phenomenon by writing f(-x) = f(x) for all x. That is, if you give x to f as an input, you get back the same answer as if you used the input -x instead. Notice that  $g(x) = x^4$  and  $h(x) = x^6$  also have this property—in fact,  $j(x) = x^n$ , where n is any even number (n could in fact be negative), has the same property. Inspired by this, we say that a function f is even if f(-x) = f(x) for all x in the domain of f. It's not good enough for this equation to be true for some values of x; it has to be true for all x in the domain of f.

Now, let's say we play the same game with  $f(x) = x^3$ . Take your favorite positive number (I'll stick with 3) and hit that with f (I get 27). Now try again with the negative of your number, -3 in my case; I get -27, and you should also get the negative of what you got before. You can express this mathematically as f(-x) = -f(x). Once again, the same property holds for  $j(x) = x^n$  when n is any odd number (and once again, n could be negative). So, we say that a function f is odd if f(-x) = -f(x) for all x in the domain of f.



In general, a function might be odd, it might be even, or it might be neither odd nor even. Don't forget this last point! Most functions are neither odd nor even. On the other hand, there's only one function that's both odd and even, which is the rather boring function given by f(x) = 0 for all x (we'll call this the "zero function"). Why is this the only odd and even function? Let's convince ourselves. If the function f is even, then f(-x) = f(x) for all x. But if it's also odd, then f(-x) = -f(x) for all x. Take the first of these equations and subtract the second from it. You should get 0 = 2f(x), which means that f(x) = 0. This is true for all x, so the function f must just be the zero function. One other nice observation is that if a function f is odd, and the number 0 is in its domain, then f(0) = 0. Why is it so? Because f(-x) = -f(x) is true for all x in the domain of f, so let's try it for x = 0. You get f(-0) = -f(0). But -0 is the same thing as 0, so we have f(0) = -f(0). This simplifies to 2f(0) = 0, or f(0) = 0 as claimed.

Anyway, starting with a function f, how can you tell if it is odd, even, or neither? And so what if it is odd or even anyway? Let's look at this second question before coming back to the first one. One nice thing about knowing that a function is odd or even is that it's easier to graph the function. In fact, if you can graph the right-hand half of the function, the left-hand half is a piece of cake! Let's say that f is an even function. Then since f(x) = f(-x), the graph of y = f(x) is at the same height above the x-coordinates x and -x. This is true for all x, so the situation looks something like this:

Section 1.4: Odd and Even Functions • 15



We can conclude that the graph of an even function has mirror symmetry about the *y*-axis. So, if you graph the right half of a function which you know is even, you can get the left half by reflecting the right half about the *y*-axis. Check the graph of  $y = x^2$  to make sure that it has this mirror symmetry.

On the other hand, let's say that f is an odd function. Since we have f(-x) = -f(x), the graph of y = f(x) is at the same height **above** the *x*-coordinate x as it is **below** the *x*-coordinate -x. (Of course, if f(x) is negative, then you have to switch the words "above" and "below.") In any case, the picture looks like this:



The symmetry is now a point symmetry about the origin. That is, the graph of an odd function has 180° point symmetry about the origin. This means that if you only have the right half of a function which you know is odd, you can get the left half as follows. Pretend that the curve is sitting on top of the paper, so you can pick it up if you like but you can't change its shape. Instead of picking it up, put a pin through the curve at the origin (remember, odd functions must pass through the origin if they are defined at 0) and then spin the whole curve around half a revolution. This is what the left-hand half of the graph looks like. (This doesn't work so well if the curve isn't continuous, that is, if the curve isn't all in one piece!) Check to see that the above graph and also the graph of  $y = x^3$  have this symmetry.



Now, suppose f is defined by the equation  $f(x) = \log_5(2x^6 - 6x^2 + 3)$ . How do you tell if f is odd, even, or neither? The technique is to calculate f(-x)by replacing every instance of x with (-x), making sure not to forget the parentheses around -x, and then simplifying the result. If you end up with the original expression f(x), then f is even; if you end up with the negative of the original expression f(-x), then f is odd; if you end up with a mess that isn't either f(x) or -f(x), then f is neither (or you didn't simplify enough!).

16 • Functions, Graphs, and Lines

In the example above, you'd write

$$f(-x) = \log_5(2(-x)^6 - 6(-x)^2 + 3) = \log_5(2x^6 - 6x^2 + 3),$$



which is actually equal to the original f(x). So the function f is even. How about

$$g(x) = \frac{2x^3 + x}{3x^2 + 5}$$
 and  $h(x) = \frac{2x^3 + x - 1}{3x^2 + 5}$ ?

Well, for g, we have

$$g(-x) = \frac{2(-x)^3 + (-x)}{3(-x)^2 + 5} = \frac{-2x^3 - x}{3x^2 + 5}.$$

Now you have to observe that you can take the minus sign out front and write

$$g(-x) = -\frac{2x^3 + x}{3x^2 + 5},$$

which, you notice, is equal to -g(x). That is, apart from the minus sign, we get the original function back. So, g is an odd function. How about h? We have

$$h(-x) = \frac{2(-x)^3 + (-x) - 1}{3(-x)^2 + 5} = \frac{-2x^3 - x - 1}{3x^2 + 5}.$$

Once again, we take out the minus sign to get

$$h(-x) = -\frac{2x^3 + x + 1}{3x^2 + 5}.$$

Hmm, this doesn't appear to be the negative of the original function, because of the +1 term in the numerator. It's not the original function either, so the function h is neither odd nor even.



Let's look at one more example. Suppose you want to prove that the product of two odd functions is always an even function. How would you go about doing this? Well, it helps to have names for things, so let's say we have two odd functions f and g. We need to look at the product of these functions, so let's call the product h. That is, we define h(x) = f(x)g(x). So, our task is to show that h is even. We'll do this by showing that h(-x) = h(x), as usual. It will be helpful to note that f(-x) = -f(x) and g(-x) = -g(x), since f and g are odd. Let's start with h(-x). Since h is the product of f and g, we have h(-x) = f(-x)g(-x). Now we use the oddness of f and g to express this last term as (-f(x))(-g(x)). The minus signs now come out front and cancel out, so this is the same thing as f(x)g(x) which of course equals h(x). We could (and should) express all this text mathematically like this:

$$h(-x) = f(-x)g(-x) = (-f(x))(-g(x)) = f(x)g(x) = h(x).$$



Anyway, since h(-x) = h(x), the function h is even. Now you should try to prove that the product of two even functions is always even, and also that the product of an odd and an even function must be odd. Go on, do it now!

Section 1.5: Graphs of Linear Functions • 7

## 1.5 Graphs of Linear Functions

Functions of the form f(x) = mx + b are called *linear*. There's a good reason for this: the graphs of these functions are lines. (As far as we're concerned, the word "line" always means "straight line.") The slope of the line is given by m. Imagine for a moment that you are in the page, climbing the line as if it were a mountain. You start at the left side of the page and head to the right, like this:



If the slope m is positive, as it is in the above picture, then you are heading uphill. The bigger m is, the steeper the climb. On the other hand, if the slope is negative, then you are heading downhill. The more negative the slope, the steeper the downhill grade. If the slope is zero, then the line is flat, or horizontal—you're going neither uphill nor downhill, just trudging along a flat line.

To sketch the graph of a linear function, you only need to identify two points on the graph. This is because there's only one line that goes through two different points. You just put your ruler on the points and draw the line. One point is easy to find, namely, the y-intercept. Set x = 0 in the equation y = mx + b, and you see that  $y = m \times 0 + b = b$ . That is, the y-intercept is equal to b, so the line goes through (0, b). To find another point, you could find the x-intercept by setting y = 0 and finding what x is. This works pretty well except in two cases. The first case is when b = 0, in which case we are just dealing with y = mx. This goes through the origin, so the x-intercept and the y-intercept are both zero. To get another point, you'll just have to substitute in x = 1 and see that y = m. So, the line y = mx goes through the origin and (1, m). For example, the line y = -2x goes through the origin and also through (1, -2), so it looks like this:



#### 18 • Functions, Graphs, and Lines

The other bad case is when m = 0. But then we just have y = b, which is a horizontal line through (0, b).

For a more interesting example, consider  $y = \frac{1}{2}x - 1$ . The *y*-intercept is -1, and the slope is  $\frac{1}{2}$ . To sketch the line, find the *x*-intercept by setting y = 0. We get  $0 = \frac{1}{2}x - 1$ , which simplifies to x = 2. So, the line looks like this:



Now, let's suppose you know that you have a line in the plane, but you don't know its equation. If you know it goes through a certain point, and you know what its slope is, then you can find the equation of the line. You really, really, really need to know how to do this, since it comes up a lot. This formula, called the *point-slope* form of a linear function, is what you need to know:



For example, what is the equation of the line through (-2, 5) which has slope -3? It is y - 5 = -3(x - (-2)), which you can expand and simplify down to y = -3x - 1.

Sometimes you don't know the slope of the line, but you do know two points that it goes through. How do you find the equation? The technique is to find the slope, then use the previous idea with one of the points (your choice) to find the equation. First, you need to know this:

If a line goes through 
$$(x_1, y_1)$$
 and  $(x_2, y_2)$ , its slope is equal to  $\frac{y_2 - y_2}{x_2 - x_2}$ 

 $\bigcirc$ 

So, what is the equation of the line through 
$$(-3, 4)$$
 and  $(2, -6)$ ? Let's find the slope first:

slope 
$$=$$
  $\frac{-6-4}{2-(-3)} = \frac{-10}{5} = -2.$ 

We now know that the line goes through (-3, 4) and has slope -2, so its equation is y - 4 = -2(x - (-3)), or after simplifying, y = -2x - 2. Alternatively, we could have used the other point (2, -6) with slope -2 to see that the equation of the line is y - (-6) = -2(x - 2), which simplifies to y = -2x - 2. Thankfully this is the same equation as before—it doesn't matter which point you pick, as long as you have used both points to find the slope.

Section 1.6: Common Functions and Graphs  $\bullet$  19

# 1.6 Common Functions and Graphs

Here are the most important functions you should know about.

**1.** Polynomials: these are functions built out of nonnegative integer powers of x. You start with the building blocks 1, x,  $x^2$ ,  $x^3$ , and so on, and you are allowed to multiply these basic functions by numbers and add a finite number of them together. For example, the polynomial  $f(x) = 5x^4 - 4x^3 + 10$  is formed by taking 5 times the building block  $x^4$ , and -4 times the building block  $x^3$ , and 10 times the building block 1, and adding them together. You might also want to include the intermediate building blocks  $x^2$  and x, but since they don't appear, you need to take 0 times of each. The amount that you multiply the building block  $x^n$  by is called the *coefficient* of  $x^n$ . For example, in the polynomial f above, the coefficient of  $x^4$  is 5, the coefficient of  $x^3$  is -4, the coefficients of  $x^2$  and x are both 0, and the coefficient of 1 is 10. (Why allow x and 1, by the way? They seem different from the other blocks, but they're not really:  $x = x^1$  and  $1 = x^0$ .) The highest number n such that  $x^n$  has a nonzero coefficient is called the *degree* of the polynomial. For example, the degree of the above polynomial f is 4, since no power of x greater than 4 is present. The mathematical way to write a general polynomial of degree n is

$$p(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_2 x^2 + a_1 x + a_0,$$

where  $a_n$  is the coefficient of  $x^n$ ,  $a_{n-1}$  is the coefficient of  $x^{n-1}$ , and so on down to  $a_0$ , which is the coefficient of 1.

Since the functions  $x^n$  are the building blocks of all polynomials, you should know what their graphs look like. The even powers mostly look similar to each other, and the same can be said for the odd powers. Here's what the graphs look like, from  $x^0$  up to  $x^7$ :



20 • Functions, Graphs, and Lines

Sketching the graphs of more general polynomials is more difficult. Even finding the x-intercepts is often impossible unless the polynomial is very simple. There is one aspect of the graph that is fairly straightforward, which is what happens at the far left and right sides of the graph. This is determined by the so-called *leading coefficient*, which is the coefficient of the highest-degree term. This is basically the number  $a_n$  defined above. For example, in our polynomial  $f(x) = 5x^4 - 4x^3 + 10$  from above, the leading coefficient is 5. In fact, it only matters whether the leading coefficient is positive or negative. It also matters whether the degree of the polynomial is odd or even; so there are four possibilities for what the edges of the graph can look like:



The wiggles in the center of these diagrams aren't relevant—they depend on the other terms of the polynomial. The diagram is just supposed to show what the graphs look like near the left and right edges. In this sense, the graph of our polynomial  $f(x) = 5x^4 - 4x^3 + 10$  looks like the leftmost picture above, since n = 4 is even and  $a_n = 5$  is positive.

Let's spend a little time on degree 2 polynomials, which are called *quadrat*ics. Instead of writing  $p(x) = a_2x^2 + a_1x + a_0$ , it's easier to write the coefficients as a, b, and c, so we have  $p(x) = ax^2 + bx + c$ . Quadratics have two, one, or zero (real) roots, depending on the sign of the *discriminant*. The discriminant, which is often written as  $\Delta$ , is given by  $\Delta = b^2 - 4ac$ . There are three possibilities. If  $\Delta > 0$ , then there are two roots; if  $\Delta = 0$ , there is one root, which is called a *double root*; and if  $\Delta < 0$ , then there are no roots. In the first two cases, the roots are given by

$$\frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$



Notice that the expression in the square root is just the discriminant. An important technique for dealing with quadratics is completing the square. Here's how it works. We'll use the example of the quadratic  $2x^2 - 3x + 10$ . The first step is to take out the leading coefficient as a factor. So our quadratic becomes  $2(x^2 - \frac{3}{2}x + 5)$ . This reduces the situation to dealing with a monic quadratic, which is a quadratic with leading coefficient equal to 1. So, let's worry about  $x^2 - \frac{3}{2}x + 5$ . The main technique now is to take the coefficient of x, which in our example is  $-\frac{3}{2}$ , divide it by 2 to get  $-\frac{3}{4}$ , and square it. We get  $\frac{9}{16}$ . We wish that the constant term were  $\frac{9}{16}$  instead of 5, so let's do some

Section 1.6: Common Functions and Graphs • 21

mental gymnastics:

$$x^{2} - \frac{3}{2}x + 5 = x^{2} - \frac{3}{2}x + \frac{9}{16} + 5 - \frac{9}{16}.$$

Why on earth would we want to add and subtract  $\frac{9}{16}$ ? Because the first three terms combine to form  $(x - \frac{3}{4})^2$ . So, we have

$$x^{2} - \frac{3}{2}x + 5 = \left(x^{2} - \frac{3}{2}x + \frac{9}{16}\right) + 5 - \frac{9}{16} = \left(x - \frac{3}{4}\right)^{2} + 5 - \frac{9}{16}$$

Now we just have to work out the last little bit, which is just arithmetic:  $5 - \frac{9}{16} = \frac{71}{16}$ . Putting it all together, and restoring the factor of 2, we have

$$2x^{2} - 3x + 10 = 2\left(x^{2} - \frac{3}{2}x + 5\right) = 2\left(\left(x - \frac{3}{4}\right)^{2} + \frac{71}{16}\right)$$
$$= 2\left(x - \frac{3}{4}\right)^{2} + \frac{71}{8}.$$

It turns out that this is a much nicer form to deal with in a number of situations. Make sure you know how to complete the square, since we'll be using this technique a lot in Chapters 18 and 19.

#### 2. Rational functions: these are functions of the form

$$\frac{p(x)}{q(x)},$$

where p and q are polynomials. Rational functions will pop up in many different contexts, and the graphs can look very different depending on the polynomials p and q. The simplest examples of rational functions are polynomials themselves, which arise when q(x) is the constant polynomial 1. The next simplest examples are the functions  $1/x^n$ , where n is a positive integer. Let's look at some of the graphs of these functions:



The odd powers look similar to each other, and the even powers look similar to each other. It's worth knowing what these graphs look like.

22 • Functions, Graphs, and Lines

**3. Exponentials and logarithms:** you need to know what graphs of exponentials look like. For example, here is  $y = 2^x$ :



The graph of  $y = b^x$  for any other base b > 1 looks similar to this. Things to notice are that the domain is the whole real line, the *y*-intercept is 1, the range is  $(0, \infty)$ , and there is a horizontal asymptote on the left at y = 0. In particular, the curve  $y = b^x$  does **not**, I repeat, not touch the *x*-axis, no matter what it looks like on your graphing calculator! (We'll be looking at asymptotes again in Chapter 3.) The graph of  $y = 2^{-x}$  is just the reflection of  $y = 2^x$  in the *y*-axis:



How about when the base is less than 1? For example, consider the graph of  $y = (\frac{1}{2})^x$ . Notice that  $(\frac{1}{2})^x = 1/2^x = 2^{-x}$ , so the above graph of  $y = 2^{-x}$  is also the graph of  $y = (\frac{1}{2})^x$ , since  $2^{-x}$  and  $(\frac{1}{2})^x$  are equal for any x. The same sort of thing happens for  $y = b^x$  for any 0 < b < 1, not just  $b = \frac{1}{2}$ .

Now, notice that the graph of  $y = 2^x$  satisfies the horizontal line test, so there is an inverse function. This is in fact the base 2 logarithm, which is written  $y = \log_2(x)$ . Using the line y = x as a mirror, the graph of  $y = \log_2(x)$  looks like this:

Section 1.6: Common Functions and Graphs • 23



The domain is  $(0, \infty)$ ; note that this backs up what I said earlier about not being able to take logarithms of a negative number or of 0. The range is all of  $(-\infty, \infty)$ , and there's a vertical asymptote at x = 0. The graphs of  $\log_{10}(x)$ , and indeed  $\log_b(x)$  for any b > 1, are very similar to this one. The log function is very important in calculus, so you should really know how to draw the above graph. We'll look at other properties of logarithms in Chapter 9.

4. Trig functions: these are so important that the entire next chapter is devoted to them.

5. Functions involving absolute values: let's take a close look at the absolute value function f given by f(x) = |x|. Here's the definition of |x|:

$$|x| = \begin{cases} x & \text{if } x \ge 0, \\ -x & \text{if } x < 0. \end{cases}$$

Another way of looking at |x| is that it is the distance between x and 0 on the number line. More generally, you should learn this nice fact:

|x - y| is the distance between x and y on the number line.



For example, suppose that you need to identify the region  $|x - 1| \leq 3$  on the number line. You can interpret the inequality as "the distance between x and 1 is less than or equal to 3." That is, we are looking for all the points that are no more than 3 units away from the number 1. So, let's take a number line and mark in the number 1 as follows:



The points which are no more than 3 units away extend to -2 on the left and 4 on the right, so the region we want looks like this:



So, the region  $|x - 1| \le 3$  can also be described as [-2, 4].

24 • Functions, Graphs, and Lines

It's also true that  $|x| = \sqrt{x^2}$ . To check this, suppose that  $x \ge 0$ ; then  $\sqrt{x^2} = x$ , no problem. If instead x < 0, then it can't be true that  $\sqrt{x^2} = x$ , since the left-hand side is positive but the right-hand side is negative. The correct equation is  $\sqrt{x^2} = -x$ ; now the right-hand side is positive, since it's minus a negative number. If you look back at the definition of |x|, you'll see that we have just proved that  $|x| = \sqrt{x^2}$ . Even so, to deal with |x|, it's much better to use the piecewise definition than to write it as  $\sqrt{x^2}$ .

Finally, let's take a look at some graphs. If you know what the graph of a function looks like, you can get the graph of the absolute value of that function by reflecting everything below the x-axis up to above the x-axis, using the x-axis as your mirror. For example, here's the graph of y = |x|, which comes from reflecting the bottom portion of y = x in the x-axis:



How about the graph of  $y = |\log_2(x)|$ ? Using the reflection of the graph of  $y = \log_2(x)$  above, this is what the absolute value version looks like:



Anyway, that's all I have to say about functions, apart from trig functions which are the subject of the next chapter. Hopefully you've seen a lot of the stuff in this chapter before. Most of the material in this chapter is used over and over again in calculus, so make sure you really get on top of it all as soon as you can!

# INDEX

absolute convergence, 491, 516 absolute convergence test for improper integrals, 447-449, 453 for series, 490-491, 516-518 absolute maximum, see global maximum absolute minimum, see global minimum absolute values, 23-24 in limits, see limits, involving absolute values acceleration, 114 constant negative, 115-117 alternating series test, 497-499, 516-518, 548antiderivatives, 361 approximations, see estimates arc lengths, 637-639 parametric formula for, 638 polar formula for, 639  $\arccos(x)$ , see inverse cosine  $\operatorname{arcsec}(x)$ , see inverse secant  $\arcsin(x)$ , see inverse sine  $\arctan(x)$ , see inverse tangent areas between curve and y-axis, 344–346 between two curves, 342-344 and definite integrals, 326 and displacement, 314-318 enclosed by polar curves, 591-593 signed, 319-320 unsigned, see unsigned areas using definite integrals to find, 339-346 argument, 601 ASTC method, 31-33 asymptotes horizontal, 47 misconceptions about, 50 vertical, 46

average velocity, 85, 350 axis, 632 base, 167 bell-shaped curve, 703 binomial theorem, 539 blow-up points, 432, see also problem spots in interior, 436 at left-hand endpoint, 433 at right-hand endpoint, 436 bounded functions, 431 cardioid, 589 Cartesian coordinates, 581, 599 and complex numbers, see Cartesian form conversion of from polar coordinates, 582 - 583conversion of to polar coordinates, 583-585Cartesian form, 600 conversion of from polar form, 601

asymptotic functions, 442, 455

average value of functions, 350

asymptotic sequences, 488

average speed, 84

conversion of to polar form, 601–603 center of power series, 529 chain rule, 107–109 justification of, 113 proof of, 693–694 change of base rule (logarithms), 171 characteristic quadratic equations, 654–656 closed interval, 3 codomain, 1 coefficients leading, 20 of polynomials, 19

of power series, 527

#### 720 • Index

coefficients (continued) of Taylor series, 530 comparison test for improper integrals, 439-441, 455 for series, 487-488, 510-515 completeness, 686, 687 completing the square, 20, 202, 402 and trig substitutions, 426 complex conjugate, 597 complex numbers, 596 adding, 596 arguments of, 601 Cartesian form of. see Cartesian form conjugates of, 597 dividing, 596-598 and exponentials, 598-599 imaginary part of, 596 modulus of, 597 multiplying, 596 polar form of, see polar form real part of, 596 representation of on complex plane, 599-603 solving  $e^{z} = w$ , 610–612 solving  $z^n = w$ , 604–610 summary of method for, 607 subtracting, 596 taking large powers of, 603-604 complex plane, 599-603 composition of functions, 11-14 compound interest, 173-175 concave down, 237 concave up, 237 conditional convergence, 498 conjugate expression, 61 constant functions, derivatives of, 102 constant multiples and derivatives, 103, 691 and integrals, 373 constant-coefficient differential equations, 653 - 665continuity on an interval, 77 at a point, 76 continuous functions, 77 compositions of, 684-686 and differentiable functions, 96-97 examples of, 77-80 convergence absolute, 491, 516 conditional, 498 of improper integrals, 433 of power series, 551-558 of sequences, 478

of series, 482 of Taylor series, 530-534 correction term, second order, 522, 523 cosecant. 27 derivative of, 143 graph of, 38 integrals involving powers of, 418 inverse of, see inverse cosecant symmetry properties of, 38  $\cosh(x)$ , see hyperbolic cosine cosine, 26 derivative of 142 graph of, 36 integrals involving powers of, 413-415 inverse of, see inverse cosine symmetry properties of, 38  $\cos(x)$ , see cosine  $\cos^{-1}(x)$ , see inverse cosine cotangent, 27 derivative of, 143 graph of, 38 integrals involving powers of, 418 inverse of, see inverse cotangent symmetry properties of, 38  $\operatorname{coth}(x)$ , see hyperbolic cotangent  $\cot(x)$ , see cotangent critical points, 227 classifying using the first derivative, 240 - 242classifying using the second derivative, 242-243  $\operatorname{csch}(x)$ , see hyperbolic cosecant  $\csc(x)$ , see cosecant cylindrical shells, see shell method decay constant, 196 decreasing functions, 236 definite integrals and areas, 326 basic idea of, 325-330 and constants, 337 definition of, 330-334 estimating, 346-350 properties of, 334-338 splitting of into two pieces, 337 and sums and differences, 338 degree of polynomial, 19 of Taylor polynomial, 533 derivatives, 90 of compositions, 107 of constant functions, 102 of constant multiples, 103, 691

of  $\cos(x)$ , 142

Index • 721

of  $\cot(x)$ , 143 of  $\csc(x)$ , 143 of differences, 103, 691-692 finding using power or Taylor series, 568 - 570higher-order, 94 implicit, see implicit differentiation of inverse functions, 204-207 involving trig functions, 141-148 left-hand, 95 as limiting ratios, 91–93 of  $\ln(x)$ , 177–179 of  $\log_{b}(x)$ , 177–179 logarithmic, 189–192 of logarithms, 177-179 nonexistence of, 94 of parametric equations, 578-580 of piecewise-defined functions, 119-123, 697-698 in polar coordinates, 590-591 of products, see product rule of quotients, see quotient rule right-hand, 95 of  $\sec(x)$ , 142 second, 94 of  $\sin(x)$ , 141 of sums, 103, 691-692 table of signs for, 247-248 of tan(x), 142 third, 94 using the definition to find, 99 using to classify critical points, 240-242using to show inverse exists, 201–203 of  $x^n$ , 101–102 difference of two cubes, 58 differentiable functions, 90 and continuous functions, 96-97 differential, 281–282 differential equations, 193, 645–646 constant-coefficient, 653-665 first order, 645 first-order homogeneous, 654 first-order linear, 648-653 and initial value problems, see initial value problems and modeling, 665-667 nonhomogeneous, 656-663 second-order homogeneous, 654-656 separable, 646–648 differentiation, 90 disc method, 619-620, 622 discontinuity, 76 discriminant, 20

displacement, 85 and areas, 314–318 as integral of velocity, 327 distance (integral of speed), 327 divergence of improper integrals, 433 of sequences, 478 of series, 482 domain, 1 finding, 4–5 restricting, 2, 9 double root, 20, 595, 656 double-angle formulas, 40, 409 dummy variable, 43, 308, 356

#### e

definition of, 173–175 limits involving, 181-182 endpoints of integration, 326 envelope, 140 equating coefficients in differential equations, 658 in partial fractions, 404 error term in linearization, 281, 285-287, 696-697 in Taylor series, 524, 536 techniques for estimating, 548–550 estimates of definite integrals, 346-350 error in, 711-714 using Simpson's rule, 709–710 using strips, 703–706 using the trapezoidal rule, 706–708 using linearization, 279–281 using quadratics, 521–522 using Taylor polynomials, 519-520, 540 - 548Euler's identity, 599, 615 even functions, 14 product of, 16 symmetry of graph of, 15 exponent, 167 exponential decay, 193, 195-197 equation describing, 197 exponential growth, 193-195 equation describing, 194 exponential rules, 168 exponentials behavior of near 0, 182-183, 472-473 behavior of near  $\pm \infty$ , 184–186, 461– 464 complex, 598-599 graph of, 22

#### 722 • Index

exponentials (continued) relationship of with logarithms, 169 theory of. 689-691 extrema, 225 Extreme Value Theorem, 227 proof of, 694-695 First Fundamental Theorem, 358–361 proof of, 381-382 solving problems using, 366-371 statement of, 360 first-order differential equations, 645 homogeneous, 654 linear, 648-653 nonhomogeneous, 656-663 form for partial fractions, 399 for particular solutions, 658, 659 functions, 1 asymptotic, 442, 455 average value of, 350 based on integral, 355–358 continuous. see continuous functions decreasing, 236 differentiable, see differentiable functions even, see even functions exponential, see exponentials hyperbolic, see hyperbolic functions increasing, 236 integrable, 331 inverse, see inverse functions inverse hyperbolic, see inverse hyperbolic functions inverse trig, see inverse trig functions involving absolute values, see absolute values linear. see linear functions logarithm, see logarithms nonintegrable, 353-354 odd, see odd functions poly-type, 67 rational, see rational functions symmetry properties of, 14-16 trigonometric, see trig functions with zero derivative, 236 Fundamental Theorem of Algebra, 595 Fundamental Theorem of Calculus First. see First Fundamental Theorem Second, see Second Fundamental Theorem geometric progressions, 480-481

geometric series, 484-485, 502-503 global maximum, 226 how to find. 228-230 global minimum, 226 how to find, 228-230 graphs of common functions, 19-24 method for sketching, 250–252 shifting, 13 growth constant, 194 half-life, 196 half-open interval, 3 harmonic series, 489 homogeneous differential equations, 654 first-order, 654 second-order, 654-656 homogeneous solutions, 658 conflicts with particular solutions, 662-663 horizontal asymptotes, 47 horizontal line test, 8-9 hyperbolic cosecant, 199 inverse of, 222–223 hyperbolic cosine, 198-200 inverse of, 220-222 hyperbolic cotangent, 199 inverse of. 222-223 hyperbolic functions, 198-200 hyperbolic geometry, 198 hyperbolic secant, 199 inverse of, 222-223 hyperbolic sine, 198–200 inverse of, 220-222 hyperbolic tangent, 199 inverse of, 222–223 imaginary numbers, 596 imaginary part, 596 implicit differentiation, 149-154 in optimization, 274-275 and second derivatives, 154–156 improper integrals, 431–476 absolute convergence test for, see absolute convergence test, for improper integrals comparison test for, see comparison test, for improper integrals definition of, 432 limit comparison test for, see limit comparison test, for improper integrals and negative function values, 453-454

Index • 723

*p*-test for, *see p*-test, for improper integrals and series, 487–491 splitting of, 452–453 summary of tests for analyzing, 454-456 increasing functions, 236 indefinite integrals, 364-366 indeterminate forms, 58, see also l'Hôpital's Bule index of summation, 308 infimum. 230 infinite sequences, see sequences infinite series, see series inflection points, 238-239 initial value problems (IVP), 646, 647 constant-coefficient linear, 663-665 input, 1 instantaneous velocity, see velocity integrable functions, 331 integral test (for series), 494-497, 509-510 integrals definite, see definite integrals improper, see improper integrals indefinite, see indefinite integrals integrand. 326 integrating factor, 649, 652-653 integration and partial fractions, see partial fractions involving powers of cos, 413-415 involving powers of cot, 418 involving powers of csc, 418 involving powers of sec, 416–418 involving powers of sin, 413–415 involving powers of tan, 415–416 involving powers of trig functions, 413-421 overview of techniques of, 429-430 by parts, 393-397 substitution method of, 383-391 using trig identities, 409–413 using trig substitutions, see trig substitutions integration by parts, 393-397 Intermediate Value Theorem (IVT), 80–82 proof of, 686-687 interval notation, 3–4 inverse cosecant derivative of. 218 domain of, 217 graph of, 217 limits at  $\pm \infty$ , 218

range of, 217 symmetry properties of, 217 inverse cosine derivative of. 212 domain of. 212 graph of, 211 range of, 212 relationship of with inverse sine, 212-213symmetry properties of, 212 inverse cotangent derivative of. 218 domain of. 217 graph of, 217 limits at  $\pm \infty$ , 218 range of, 217 symmetry properties of, 217 inverse functions, 7-8 derivatives of, 204–207 existence of, 201-203 finding, 9 inverses of, 11 inverse hyperbolic cosecant, 222-223 inverse hyperbolic cosine, 220–222 inverse hyperbolic cotangent, 222-223 inverse hyperbolic functions, 220–223 inverse hyperbolic secant, 222–223 inverse hyperbolic sine, 220–222 inverse hyperbolic tangent, 222–223 inverse secant derivative of, 217 domain of, 217 graph of, 216 limits at  $\pm \infty$ , 216 range of, 217 symmetry properties of, 217 inverse sine, 208–211 derivative of, 210 domain of, 210 graph of, 209 range of, 210 relationship of with inverse cosine, 212-213symmetry properties of, 210 inverse tangent derivative of, 215 domain of, 215 graph of, 214 limits at  $\pm \infty$ , 215 range of, 215 symmetry properties of, 215 inverse trig functions, 208–218 computing, 218-220

#### $724 \bullet Index$

IVP, see initial value problems IVT, see Intermediate Value Theorem large numbers, 48–49 leading coefficient, 20 left-continuous functions, 77 left-hand derivatives. see derivatives. lefthand left-hand limits, see limits, left-hand l'Hôpital's Rule, 293–303 proof of, 698-700 for sequences, 479 summary of, 302-303 Type A (0/0), 294–296 Type **A**  $(\infty/\infty)$ , 296–297 Type **B1** ( $\infty - \infty$ ), 298–299 Type **B2**  $(0 \times \pm \infty)$ , 299–300 Type **C**  $(1^{\pm \infty}, 0^0, \text{ or } \infty^0), 301-302$ limacon, 589 limit comparison test for improper integrals, 441-444, 455-456for series, 488-489, 510-515 limits as derivative in disguise, 117-119 finding using Maclaurin series, 570-574formal definition of, 672 game about, 670-672 infinite, 45, 679-680 at  $\infty$  or  $-\infty$ , 47, 680–682 informal definition of, 42 involving absolute values, 72–73 involving definition of e, 181–182 involving poly-type functions, 66-70 involving rational functions as  $x \to a$ , 57–60 as  $x \to -\infty$ , 70–72 as  $x \to \infty$ , 61–66 involving square roots, 61 involving trig functions, 127–141 at large arguments, 134–136 at small arguments, 128–133 at various other arguments, 137 left-hand, 43–44, 680 nonexistence of, 44-46 overview of methods involving, 303-306 products of, 675-676 quotients of, 676-677 right-hand, 43-44, 680 of sequences, 682 summary of basic types of, 54

sums and differences of, 674-675 two-sided, 44 limits of integration, 326 equal, 335 reversing, 335 linear functions, 17 derivatives of, 93 graph of, 17 point-slope form of, 18 through two given points, 18 linearization, 278-281, 520-521 error in. see error term. in linearization method for finding, 283-285  $\ln(x)$ , see logarithms local maximum, 226 local minimum, 226 log, see logarithms log rules, 171-172, 176 logarithmic differentiation, 189–192 logarithms behavior of near 0, 188-189, 473-474 behavior of near 1, 183-184 behavior of near  $\infty$ , 187–188, 465– 468 definition of, 168 derivatives of, 177-179 graph of, 22 natural, 176 relationship of with exponentials, 169 rules for, see log rules theory of, 689-691 lower sums, 324, 354 Maclaurin series, 529-530, 536, 615 common, 558-559

and improper integrals, 474-475 using to find limits, 570-574 Max-Min Theorem, 83 proof of, 687-688 maximum, 83 global, see global maximum local, see local maximum Mean Value Theorem (MVT), 233–235, 525 consequences of, 235–236 for integrals, 351-353 proof of, 695-696 mesh, 322, 330 minimum, 83 global, see global minimum local, see local minimum mod-arg form, 601 modulus, 597

Index • 725

monic quadratics, 20 MVT, see Mean Value Theorem natural logarithms, 176 Newton's method, 287–292 formula for, 289 potential problems with, 290–292 nonhomogeneous differential equations, 656-663 nonintegrable functions, 353-354 nonnegative numbers, 2 normal distribution, 703 nth term test, 486-487, 503-504 odd functions, 14 product of, 16 symmetry of graph of, 15 open interval, 3 optimization, 267-278 method for solving problems involving, 269 using implicit differentiation in, 274-275order of differential equations, 645 of Taylor polynomials, 533 output, 1 overestimates in linearization, 286 in Taylor series, 541 p-test for improper integrals, 456 for improper integrals, 444–447 for series, 489-490, 497, 510-515 parameters, 576 and arc lengths, 638 and surface areas, 643 parametric equations, 575–578 derivatives of, 578-580 second derivatives of, 580-581 parametrization, 577 and speed, 639-640 partial fractions, 397–408 form for, 399 main method of, 404 partial sums, 482, 483 particular solutions, 657 conflicts with homogeneous solutions, 662 - 663finding, 658-662 partitions, 317, 330, 704 evenly spaced, 705–706 mesh of, see mesh

parts, integration by, 393-397 periodic, 35, 589, 601 piecewise-defined functions, derivatives of, 697-698 point-slope form, 18 points of inflection, 238-239 polar coordinates, 581-590, 599 and arc lengths, 639 and complex numbers. see polar form conversion of from Cartesian coordinates. 583-585 conversion of to Cartesian coordinates. 582 - 583sketching curves in, 585–590 polar curves areas enclosed by, 591-593 tangents to, 590-591 polar form, 600 conversion of from Cartesian form, 601-603 conversion of to Cartesian form, 601 poly-type functions, 67 behavior of near 0, 469-470 behavior of near  $\pm \infty$ , 456–459 in limits, see limits, involving polytype functions polynomials, 19 behavior of near 0, 469-470 behavior of near  $\pm \infty$ , 456–459 coefficients of, 19 degree of, 19 leading coefficient of, 20 power series, 527-529, 615 convergence of, 551-558 radius of convergence of, 551-558 using to find derivatives, 568-570 powers of x, derivatives of, 101–102, 192– 193problem spots, 436, 437, 451 absence of, 452 not at 0 or  $\infty$ , 475–476 product rule, 104-105 for three variables, 112 justification of, 111–113 proof of, 692 for three variables, 105 quadrant, 28 quadratics, 20-21 completing the square in, 20 and complex numbers, 598 discriminant of, 20 double root of, 20

#### 726 • Index

quadratics (continued) monic, 20 quotient rule, 105-106 proof of, 693 radar. 584 radians. 25 radius of convergence, 551-558 range, 2 finding, 5-6 rates of change, 156 ratio test, 492-493, 504-508 rational functions, 21 in limits, see limits, involving rational functions integrating, see partial fractions real part, 596 reduction formulas. 419-421 reference angle, 30 related rates, 156-165 relative maximum, see local maximum relative minimum, see local minimum remainder term, see error term, in Taylor series restricting the domain, see domain, restricting Riemann sums, 331, 333, 355, 591, 618, 704 right-continuous functions, 77 right-hand derivatives, see derivatives, righthand right-hand limits, see limits, right-hand Rolle's Theorem, 230-233 proof of, 695 root test, 493-494, 508 sandwich principle, 51–54 proof of, 678 for sequences, 479 secant, 27 derivative of, 142 graph of, 37 integrals involving powers of, 416-418 inverse of. see inverse secant symmetry properties of, 38  $\operatorname{sech}(x)$ , see hyperbolic secant second derivatives, 94 and graphs, 237-239 and implicit differentiation, 154-156 of parametric equations, 580-581 table of signs for, 248-250 using to classify critical points, 242 Second Fundamental Theorem, 362–364 solving problems using, 371-374

statement of, 363 second-order correction term, 522, 523 second-order differential equations homogeneous, 654-656 nonhomogeneous, 656-663  $\sec(x)$ , see secant  $\sec^{-1}(x)$ , see inverse secant separable differential equations, 646-648 sequences, 477 asymptotic, 488 and functions, 478-480 limits of 682 series absolute convergence of, 491, 516 absolute convergence test for, see absolute convergence test, for series alternating series test for, see alternating series test basic concepts for, 481-484 comparison test for, see comparison test, for series conditional convergence of, 498 flowchart for investigating, 501–502 geometric, *see* geometric series harmonic, 489 and improper integrals, 487–491 integral test for, see integral test (for series) limit comparison test for, see limit comparison test, for series Maclaurin. see Maclaurin series with negative terms, 515–518 nth term test for, see nth term test *p*-test for, *see p*-test, for series power, see power series ratio test for, see ratio test root test for, see root test Taylor, see Taylor series telescoping, 311-314 shell method, 620-622 sigma notation, 307-314 signed areas, 319-320 simple harmonic motion, 145–146 Simpson's rule, 709–710 error in, 711-714 proof of, 710-711 sine, 26 derivative of. 141 graph of, 35 important limit involving, 137-140 integrals involving powers of, 413-415 inverse of, see inverse sine symmetry properties of, 38

Index • 727

 $\sinh(x)$ , see hyperbolic sine  $\sin(x)$ , see sine  $\sin^{-1}(x)$ , see inverse sine sketching graphs, 250–266 of derivatives, 123–126 in polar coordinates, 585–590 slicing, 620, 632 small numbers, 48–49 smoothness, 75 solids of revolution surface areas of, see surface areas, of solids of revolution volumes of, see volumes, of solids of revolution speed average, 84 and parametrization, 639-640 spiral of Archimedes, 589 squeeze principle, see sandwich principle standard form, of first-order differential equation, 650 strips, estimating integrals using, 703-706 substitution (integration technique), 383-391justification of, 392-393 surface areas of solids of revolution, 640– 644 parametric formula for, 643 table of signs, 245-247 for the derivative, 247-248 for the second derivative, 248-250 tangent (function), 26 derivative of, 142 graph of, 37 integrals involving powers of, 415–416 inverse of, see inverse tangent symmetry properties of, 38 tangent line, 88–90, see also linearization finding equation of, 114 tanh(x), see hyperbolic tangent  $\tan(x)$ , see tangent (function)  $\tan^{-1}(x)$ , see inverse tangent Taylor approximation theorem, 522 proof of, 700-702 Taylor polynomials, 522, 535-536 finding, 537-539 Taylor series, 529-530, 535-536 adding, 565-566 convergence of, 530-534 differentiating, 562–563 dividing, 567-568 error term in, see error term, in Taylor series

finding, 537-539 getting new from old, 558-568 integrating, 563–565 multiplying, 566-567 remainder term in, see error term, in Taylor series and substitution, 560-561 subtracting, 565-566 using to find derivatives, 568–570 Taylor's Theorem, 523–526 telescoping series, 311-314 third derivatives, 94 trapezoidal rule, 706-708 error in, 711-714 triangle inequality, 674 trig functions basic properties of, 25 behavior of near 0, 470-471 behavior of near  $\pm \infty$ , 459–461 derivatives involving, see derivatives, involving trig functions extending the domain of, 28-35 graphs of, 35-38 integrals involving powers of, 413–421 limits involving, see limits, involving trig functions periodicity of, 35 symmetry properties of, 38 trig identities complementary, 39 integration involving, 409-413 involving double angles, 40 involving sums and differences, 40 Pythagorean, 39, 410 trig substitutions, 421–429 and completing the square, 426 and square roots, 427-429 summary of, 426-427 trigonometric series, 612–615 triple-boxed principle, 605 unbounded region of integration, 437 underestimates in linearization, 286 in Taylor series, 541 unit circle, 30 unsigned areas and absolute values, 376-379 finding using definite integrals, 339-342upper sums, 324, 333, 354 velocity, 86-87, 114 average, 85, 350

#### 728 • Index

velocity (continued) continuous, 320–323 and derivatives, 91 graphical interpretation of, 87 vertical asymptotes, 46 vertical line test, 6–7 volumes of general solids, 631–637 of generalized cones, 632–636 by slicing, 620, 632 of solids of revolution, 617–631 disc method for finding, *see* disc method

of regions between curve and y-axis, 623–624

of regions between two curves, 625-628

shell method for finding, *see* shell method

whoop-di-doo, 440, 455