

CONTENTS

	<i>Preface to the Princeton Science Library Edition</i>	ix
	<i>Preface</i>	xiii
	Introduction	1
CHAPTER 1	Leonhard Euler and His Three “Great” Friends	10
CHAPTER 2	What Is a Polyhedron?	27
CHAPTER 3	The Five Perfect Bodies	31
CHAPTER 4	The Pythagorean Brotherhood and Plato’s Atomic Theory	36
CHAPTER 5	Euclid and His <i>Elements</i>	44
CHAPTER 6	Kepler’s Polyhedral Universe	51
CHAPTER 7	Euler’s Gem	63
CHAPTER 8	Platonic Solids, Golf Balls, Fullerenes, and Geodesic Domes	75
CHAPTER 9	Scooped by Descartes?	81
CHAPTER 10	Legendre Gets It Right	87
CHAPTER 11	A Stroll through Königsberg	100
CHAPTER 12	Cauchy’s Flattened Polyhedra	112
CHAPTER 13	Planar Graphs, Geoboards, and Brussels Sprouts	119

CHAPTER 14	It's a Colorful World	130
CHAPTER 15	New Problems and New Proofs	145
CHAPTER 16	Rubber Sheets, Hollow Doughnuts, and Crazy Bottles	156
CHAPTER 17	Are They the Same, or Are They Different?	173
CHAPTER 18	A Knotty Problem	186
CHAPTER 19	Combing the Hair on a Coconut	202
CHAPTER 20	When Topology Controls Geometry	219
CHAPTER 21	The Topology of Curvy Surfaces	231
CHAPTER 22	Navigating in n Dimensions	241
CHAPTER 23	Henri Poincaré and the Ascendance of Topology	253
EPILOGUE	The Million-Dollar Question	265
<i>Acknowledgments</i>		271
<i>Appendix A</i>	<i>Build Your Own Polyhedra and Surfaces</i>	273
<i>Appendix B</i>	<i>Recommended Readings</i>	283
<i>Notes</i>		287
<i>References</i>		295
<i>Illustration Credits</i>		309
<i>Index</i>		311

INTRODUCTION

Philosophy is written in this grand book—I mean the universe—which stands continually open to our gaze, but it cannot be understood unless one first learns to comprehend the language and interpret the characters in which it is written. It is written in the language of mathematics, and its characters are triangles, circles, and other geometrical figures, without which it is humanly impossible to understand a single word of it; without these one is wandering about in a dark labyrinth.

—Galileo Galilei¹

They all missed it. The ancient Greeks—mathematical luminaries such as Pythagoras, Theaetetus, Plato, Euclid, and Archimedes, who were infatuated with polyhedra—missed it. Johannes Kepler, the great astronomer, so in awe of the beauty of polyhedra that he based an early model of the solar system on them, missed it. In his investigation of polyhedra the mathematician and philosopher René Descartes was but a few logical steps away from discovering it, yet he too missed it. These mathematicians, and so many others, missed a relationship that is so simple that it can be explained to any schoolchild, yet is so fundamental that it is part of the fabric of modern mathematics.

The great Swiss mathematician Leonhard Euler (1707–1783)—whose surname is pronounced “oiler”—did not miss it. On November 14, 1750, in a letter to his friend, the number theorist Christian Goldbach (1690–1764), Euler wrote, “It astonishes me that these general properties of stereometry [solid geometry] have not, as far as I know, been noticed by anyone else.”² In this letter Euler described his observation, and a year later he gave a proof. This observation is so basic and vital that it now bears the name *Euler’s polyhedron formula*.

A polyhedron is a three-dimensional object such as those found in figure I.1. It is composed of flat polygonal *faces*. Each pair of adjacent faces meets along a line segment, called an *edge*, and adjacent edges meet at a corner, or a *vertex*. Euler observed that the numbers of vertices, edges,




Figure I.1. A cube and a soccer ball (truncated icosahedron) both satisfy Euler's formula.

and faces (V , E , and F) always satisfy a simple and elegant arithmetic relationship:

$$V - E + F = 2.$$

The cube is probably the best-known polyhedron. A quick count shows that it has six faces: a square on the top, a square on the bottom, and four squares on the sides. The boundaries of these squares form the edges. Counting them, we find twelve total: four on the top, four on the bottom, and four vertical edges on the sides. The four top corners and the four bottom corners are the eight vertices of the cube. Thus for the cube, $V = 8$, $E = 12$, and $F = 6$, and of course

$$8 - 12 + 6 = 2,$$

as claimed. It is more tedious to count them, but the soccer-ball-shaped polyhedron shown in figure I.1 has 32 faces (12 pentagons and 20 hexagons), 90 edges, and 60 vertices. Again,

$$60 - 90 + 32 = 2.$$

In addition to his work with polyhedra, Euler created the field of *analysis situs*, known today as topology. Geometry is the study of rigid objects. Geometers are interested in measuring quantities such as area, angle, volume, and length. Topology, which has inherited the popular moniker "rubber-sheet geometry," is the study of malleable shapes. The objects a topologist studies need not be rigid or geometric. Topologists are interested in determining connectedness, detecting holes, and investigating twistedness. When a carnival clown bends a balloon into the shape of a dog, the balloon is still the same topological entity, but geometrically it is very different. But when a child bursts the balloon with a pencil, leaving a gaping




Figure I.2. Topological surfaces: a sphere, a torus, and a Möbius band.




Figure I.3. Two partitions of the sphere.

hole in the rubber, it is no longer topologically the same. In figure I.2 we see three examples of topological surfaces—the sphere, the doughnut-shaped torus, and the twisted Möbius band.

Scholars in this young field of topology were fascinated by Euler’s formula, and they attempted to apply it to topological surfaces. The obvious question arose: where are the vertices, edges, and faces on a topological surface? The topologists disregarded the rigid rules set forth by the geometers and allowed the faces and edges to be curved. In figure I.3 we see a partition of a sphere into “rectangular” and “triangular” regions. The partition is formed by drawing 12 lines of longitude, meeting at the two poles, and 7 lines of latitude. This globe has 72 curved rectangular faces and 24 curved triangular ones (the triangular faces are located near the north and south poles), giving a total of 96 faces. There are 180 edges and 86 vertices. Thus, as with polyhedra, we find that

$$V - E + F = 86 - 180 + 96 = 2.$$

Likewise, the 2006 World Cup soccer ball, which consists of six four-sided hourglass-shaped patches and eight misshapen hexagonal patches




Figure I.4. A partition of the torus.

(see figure I.3), also satisfies Euler's formula (it has $V = 24$, $E = 36$, and $F = 14$).

At this point we are tempted to conjecture that Euler's formula applies to every topological surface. However, if we partition a torus into rectangular faces, as in figure I.4, we obtain a surprising result. This partition is formed by placing 2 circles around the central hole of the torus and 4 circles around its circular tube. The partition has 8 four-sided faces, 16 edges, and 8 vertices. Applying Euler's formula we find

$$V - E + F = 8 - 16 + 8 = 0,$$

rather than the expected 2.

If we were to construct a different partition of the torus we would find that the alternating sum is still zero. This gives us a *new* Euler's formula for the torus:

$$V - E + F = 0.$$

We can prove that every topological surface has its "own" Euler's formula. No matter whether we partition the surface of a sphere into 6 faces or 1,006 faces, when we apply Euler's formula we will always get 2. Likewise, if we apply Euler's formula to any partition of the torus, we will get 0. This special number can be used to distinguish surfaces just as the number of wheels can be used to distinguish highway vehicles. Every car has four wheels, every tractor trailer has eighteen wheels, and every motorcycle has two wheels. If a vehicle does not have four wheels, then it is not a car; if it does not have two wheels, then it is not a motorcycle. In the same way, if $V - E + F$ is not 0, then topologically the surface is not a torus.

The sum $V - E + F$ is a quantity intrinsically associated with the shape. In the lingo of topologists, we say that it is an *invariant* of the surface. Because of this powerful property of invariance, we call the number




Figure I.5. Are these the same knot?

$V - E + F$ the *Euler number* of the surface. The Euler number of a sphere is 2 and the Euler number of a torus is 0.

At this point, the fact that every surface has its own Euler number may seem nothing more than a mathematical curiosity, an “isn’t that cool” kind of fact to contemplate while holding a soccer ball or looking at a geodesic dome. This is most certainly not the case. As we will see, the Euler number is an indispensable tool in the study of polyhedra, not to mention topology, geometry, graph theory, and dynamical systems, and it has some very elegant and unexpected applications.

A mathematical knot is like an intertwined loop of string as shown in figure I.5. Two knots are the same if one knot can be deformed into the other without cutting and re-gluing the string. Just as we can use the Euler number to help distinguish two surfaces, with a little ingenuity we can also use it to distinguish knots. We can use the Euler number to prove that the two knots in figure I.5 are not the same.

In figure I.6 we see a snapshot of wind patterns on the surface of the earth. In this example there is a point off the coast of Chile where the wind is not blowing. It is located in the calm spot within the eye of the storm that is spinning clockwise. We can prove that there is always at least one point on the surface of the earth where there is no wind. This follows not from an understanding of meteorology, but from an understanding of topology. The existence of this point of calm comes from a theorem that mathematicians refer to as the hairy ball theorem. If we think of the wind directions as strands of hair on the surface of the earth, then there must be some point where the hair forms a cowlick. Colloquially, we say that “you can’t comb the hair on a coconut.” In chapter 19 we will see how the Euler number enables us to establish this bold assertion.



Figure I.6. Is there always a windless location on earth?




Figure I.7. Is it possible to determine the area of the shaded polygon by counting dots?

In figure I.7 we see a polygon sitting in an array of dots spaced one unit apart. The vertices of the polygon are located at the dots. Surprisingly, we can compute the precise area of the polygon simply by counting dots. In chapter 13 we will use the Euler number to derive the following elegant formula that gives the area of the polygon in terms of the number of dots that lie along the boundary of the polygon (B) and the number of dots in the interior of the polygon (I):

$$\text{Area} = I + B/2 - 1.$$

From this formula we conclude that our polygon's area is $5 + 10/2 - 1 = 9$.

There is an old and interesting problem that asks how many colors are required to color a map in such a way that every pair of regions with a common border are not the same color. Take a blank map of the United States and color it with as few crayons as possible. You will quickly discover that most of the country can be colored using only three crayons, but that a




Figure I.8. Can we color the map of the United States using only four colors?




Figure I.9. The C_{60} buckminsterfullerene molecule.

fourth is needed to complete the map. For instance, since an odd number of states surround Nevada, you will need three crayons to color them—then you will need that fourth crayon for Nevada itself (figure I.8). If we are clever, then we can finish the coloring without using a fifth—four colors are sufficient for the entire map of the United States. It was long conjectured that every map can be colored with four or fewer colors. This infamous and very slippery conjecture became known as the four color problem. In chapter 14 we will recount its fascinating history, one that ended with a controversial proof in 1976 in which the Euler number played a key role.

Graphite and diamond are two materials whose chemical makeup consists entirely of carbon atoms. In 1985 three scientists—Robert Curl Jr., Richard Smalley, and Harold Kroto—shocked the scientific community by discovering a new class of all-carbon molecules. They called these molecules *fullerenes*, after the architect Buckminster Fuller, inventor of the geodesic dome (figure I.9). They chose this name because fullerenes are large




Figure I.10. The five regular solids.

polyhedral molecules that resemble such structures. For the discovery of fullerenes the three men were awarded the 1996 Nobel Prize in chemistry. In a fullerene every carbon atom bonds with exactly three of its neighbors, and rings of carbon atoms form pentagons and hexagons. Initially Curl, Smalley, and Kroto found fullerenes possessing 60 and 70 carbon atoms, but other fullerenes were discovered later. The most plentiful fullerene is the soccer-ball-shaped molecule C_{60} that they called the buckminsterfullerene. Remarkably, knowing no chemistry and only Euler's formula, we are able to conclude that there are certain configurations of carbon atoms that are impossible in a fullerene. For instance, every fullerene, regardless of the size, must have exactly 12 pentagonal carbon rings even though the number of hexagonal rings can vary.

For thousands of years people have been drawn to the beautiful and alluring regular solids—polyhedra whose faces are identical regular polygons (figure I.10). The Greeks discovered these objects, Plato incorporated them into his atomic theory, and Kepler based an early model of the solar system on them. Part of the mystery surrounding these five polyhedra is that they are so few in number—no polyhedron other than these five satisfies the strict requirements of regularity. One of the most elegant applications of Euler's formula is a very short proof that guarantees that only five regular solids exist.

Despite the importance and beauty of Euler's formula, it is virtually unknown to the general public. It is not found in the standard curriculum taught in the schools. Some high school students may know Euler's formula, but most students of mathematics do not encounter this relation until college.

Mathematical fame is a curious thing. Some theorems are well known because they are drilled into the heads of young students: the Pythagorean theorem, the quadratic formula, the fundamental theorem of calculus. Other results are thrust into the spotlight because they resolve a famous unsolved problem. Fermat's last theorem remained unproved for over three hundred years until Andrew Wiles surprised the world with his proof in

1993. The four color problem was posed in 1853 and was only proved by Kenneth Appel and Wolfgang Haken in 1976. The famous Poincaré conjecture was posed in 1904 and was one of the Clay Mathematics Institute Millennium Problems—a collection of seven problems deemed so important that the mathematician who solves one receives \$1 million. The money is likely to be awarded to Grisha Perelman, who gave a proof of the Poincaré conjecture in 2002. Other mathematical facts are well known because of their cross-disciplinary appeal (the Fibonacci sequence in nature) or their historical significance (the infinitude of primes, the irrationality of π).

Euler's formula should be as well known as these great theorems. It has a colorful history, and many of the world's greatest mathematicians contributed to the theory. It is a deep theorem, and one's appreciation for this depth grows with one's mathematical sophistication.

This is the story of Euler's beautiful theorem. We will trace its history and show how it formed a bridge from the polyhedra of the Greeks to the modern field of topology. We will present many surprising guises of Euler's formula in geometry, topology, and dynamical systems. We will also give examples of theorems whose proofs rely on Euler's formula. We will see why this long-unnoticed formula became one of the most beloved theorems in mathematics.

INDEX

- 180° theorem, 221, 227
- Académie des Sciences* (French Academy of Sciences), 12, 14, 21, 112, 211
- Académie Royale des Sciences et Belles Lettres*. *See* Berlin Academy of Sciences
- Academy of Sciences in St. Petersburg, 14–17, 23, 103
- acies*, 64, 66
- adjacency graph, 135–137, 140–142
- Afghan bands, 163, 164, 176
- Aleksandrov, Pavel, 263
- Alekseyevna, Catherine. *See* Catherine II of Russia
- alternating knot, 194, 195, 197, 199, 200
- American Mathematical Society, 142, 158
- Anderson, Gary, 163
- angle defect. *See* angle deficit
- angle deficit: geodesic polygon, 227, 228, 236; polygon, 221, 223; polyhedron, 223–226, 228, 229
- angle excess: geodesic polygon, 227–230, 236–239; polyhedron, 224, 225, 228, 229
- angle excess theorem, 219, 223, 229, 230, 238
- angulus solidus*, 64, 66
- annulus. *See* cylinder
- Apollonius, 31, 50
- Appel, Kenneth, 8, 142–144
- Arago, François, 17
- Archimedean solid. *See* semiregular polyhedron
- Archimedes, 1, 10, 31, 48–50, 181
- Aristotle, 43, 51, 58
- ball (B^n), 216–218, 244, 247
- Ball, W. W. Rouse, 104
- Baltzer, Richard, 133
- band, 191–194, 196
- Barbari, Jacopo de, 51
- Barbaro, Daniele, 51
- Barr, Stephen, 139, 164, 284
- Berlin Academy of Sciences, 14, 18–22
- Bernoulli, Daniel, 15, 16
- Bernoulli, Jacob, 12
- Bernoulli, Johann, 12, 15
- Bernoulli, Nicolaus, 15
- Berthollet, Claude Louis, 112
- Betti, Enrico, 250–252, 254, 255, 258
- Betti group, 263
- Betti number, 255–261, 263, 266
- Biggs, Henry, 92
- Billingsley, Henry, 27
- Birkhoff, George D., 141
- Biron, Earnst-Johann, 16, 18
- Blaschke, Wilhelm, 239
- Bonnet, Pierre Ossian, 238, 264
- boundary. *See* surface with boundary; manifold with boundary
- Boyer, Carl, 25
- Boyle, Robert, *The Sceptical Chymist*, 43, 58
- Brahe, Tycho, 57
- bridges of Königsberg problem, 100–111, 135, 156, 187, 253, 269, 284
- Brouwer fixed point theorem, 216–218, 285
- Brouwer, Luitzen Egbertus Jan, 209, 212, 213, 217, 246, 263
- Brussels sprouts, 128, 129
- buckminsterfullerene, 7, 8, 78
- Burkert, Walter, 36
- calculus, ix, 14, 25, 26, 52, 113, 133, 183, 203, 231, 232
- Carroll, Lewis, 130
- Catherine I of Russia, 14, 15
- Catherine II of Russia, 11, 14, 22–24
- Catherine the Great. *See* Catherine II of Russia
- Cauchy, Augustin-Louis, x, 48, 112–119, 121, 145, 151, 152, 246, 247, 259
- Cauchy's rigidity theorem, 48, 114, 284
- Cayley, Arthur, 118, 132, 137, 156
- center (zero of a vector field), 205, 206, 208
- Challenger, HMS, 33
- Chanut, Hector-Pierre, 83
- Charles X of France, 113
- Christina, Queen of Sweden, 83

- Clarke, Arthur C., "The Wall of Darkness," 163
- classification of surfaces, 181, 183–185, 187, 246, 255, 265, 285
- Clay Mathematics Institute, 9, 268, 270
- Clerseilier, Claude, 83
- Collegium Carolinum, 234
- colorability, knot, 197, 198, 200
- compact surface, 159, 178
- complete bipartite graph, 122–124
- complete graph, 122–124, 134
- composite knot, 196
- connectivity number, 250–252, 254, 257
- Connelly, Robert, 48
- convexity, 29, 33, 46–49, 71–73, 95, 97–99, 114–117, 125, 145, 146, 148, 150–152, 155, 174, 220, 224, 225, 246, 247, 249
- Conway, John Horton, 127, 128, 184, 270; ZIP proof, 184
- Copernican astronomy, 55–57
- Copernicus, Nicolaus, 55
- Coxeter, Harold Scott MacDonald, 133, 141
- Critias, 42
- cross cap, 171, 172, 180–182, 184, 242, 246, 258
- crossing number, 198–200
- Crowell, Richard Henry, 195
- cube, 2, 27, 28, 32–34, 36, 40, 42, 46, 47, 55, 58–61, 64, 67–69, 75, 77, 78, 84, 117, 147, 153, 174, 223–225, 241, 245–247
- Curl, Robert Jr., 7, 8
- curvature; Gaussian, 233, 235–239; plane curve, 231–233; principle, 232, 233, 235, 237
- cycle, 255–259, 263
- cylinder, 154, 158–162, 167, 169, 171, 175–177, 179, 181, 182, 184, 235–237, 249, 250, 255, 273
- Czartoryski, Prince Adam, 146
- D'Alembert, Jean, 21
- de Careil, Foucher, 83
- De Morgan, Augustus, 132
- Dehn, Max Wilhelm, 184
- Descartes, René, x, 1, 62, 81–86, 224, 225, 264, 284; *Discourse on the Method for Rightly Directing One's Reason and Searching for Truth in the Sciences*, 82; *The Elements of Solids*, 83, 224
- Descartes' formula, 84, 219, 221, 223, 225, 226, 228–230
- Descartes-Euler formula. *See* Euler's polyhedron formula
- Diderot, Denis, 22
- Dieudonné, Jean Alexandre Eugène, 212, 253
- differential equation, 10, 113, 183, 202, 203, 212
- Diophantus, 50
- dipole (zero of a vector field), 205, 206, 208
- disk, 158, 159, 169, 171, 178, 179, 181, 182, 184, 185, 189–194, 196, 206, 207, 214, 217, 236, 249, 250, 255
- dodecahedron, 32–34, 36, 39, 40, 42, 46, 47, 56, 58–61, 76–78, 260, 266
- duality. *See* Poincaré duality; regular polyhedron, duality
- Dürer, Albrecht, 51
- Dyck, Walther von, 165, 166, 184, 239, 244
- Dyck's surface, 184
- dynamical systems, x, xii, 5, 9, 203, 212, 216, 267
- École Polytechnique, 112, 113
- Ehler, Carl, 101
- Einstein, Albert, 201
- Elizabeth I of Russia, 18, 21
- Empedocles, 42
- equilibrium, 203, 204
- Erdős, Paul, 144
- Erlangen University, 262
- Escher, Maurits Cornelis, 163
- Euclid, x, 1, 27, 31, 32, 44–48, 50, 51, 75, 87, 89, 123; *Elements*, 27, 32, 40, 41, 44–47, 75, 87, 89
- Eudoxus, 44
- Euler, Leonhard, x, 1, 10–27, 29, 48, 62, 81, 84–87, 91, 97, 99–107, 112, 113, 116, 118, 145, 146, 152, 156, 180, 211, 225, 233, 234, 250, 269, 270, 283; "Demonstratio nonnullarum insignium proprietatum quibus solida hedris planis insunt praedita," 66; "Elementa doctrinae solidorum," 66
- Euler, Marguerite Brucker, 11
- Euler, Paul, 11, 12
- Euler characteristic. *See* Euler number

- Euler circuit. *See* Euler walk
- Euler number, 4–7, 173, 177–181, 184, 185, 187, 189, 193, 201, 205, 208–210, 212, 213, 215, 216, 218, 219, 225, 227, 229, 230, 239–241, 247, 251, 259, 265, 284
- Euler-Poincaré characteristic, 259–263, 266
- Euler walk, 104–111
- Euler’s polyhedron formula, x–xii, 1–4, 8, 9, 26, 28, 29, 32, 46, 58, 75–77, 79, 81, 84–87, 94–101, 112, 114–125, 127, 129, 136, 145–149, 151, 152, 155–157, 174, 179, 241, 246, 247, 249, 250, 269, 270, 283–285
- Eulerian polyhedron, 146, 148, 152, 173, 174
- exterior angle theorem, 220–223, 225, 232
- Ferdinand, Carl Wilhelm, Duke of Brunswick, 234, 235
- Fermat, Pierre de, 8, 26, 82
- Fermat’s last theorem, 8, 267, 269
- Fibonacci sequence, 9
- Fields Medal, 265, 269, 270
- figure eight knot. *See* knot, figure eight
- five color theorem, 139–141
- five neighbors theorem, 136, 137, 140, 141
- fixed point, 204, 205, 208, 216–218
- flow, 203–205, 208, 216, 218; gradient, 205, 206
- four classical elements, 42, 43, 58, 59
- four color theorem, 7, 8, 130–144, 267, 284, 285
- Francesca, Piero della, 51
- Frederick II of Prussia, 11, 18–23
- Frederick the Great. *See* Frederick II of Prussia
- Frederick William I of Prussia, 18
- Freedman, Michael, 268, 269
- Freudenthal, Hans, 113
- Fuller, Buckminster, 7
- fullerene, 7, 8, 78
- Galilei, Galileo, 56
- Gardner, Martin, 127, 133, 135
- Gattegno, Caleb, 125
- Gauss, Carl Friedrich, x, 10, 25, 75, 133, 152, 181, 182, 188, 211, 231, 233–235, 237–239, 248, 264, 270
- Gauss, Gerhard, 233
- Gauss-Bonnet theorem: global, 219, 239; local, 219, 238, 239
- genus: of a knot, 194–200; of a surface, 169, 170, 181, 183, 184, 193, 194, 239, 251
- geoboard, 125, 126
- geocentric astronomy. *See* Ptolemaic astronomy
- geodesic, 89, 227–229, 238. *See also* geodesic polygon
- geodesic dome, 5, 7, 80
- geodesic polygon, 89–91, 93–96, 156, 227–229, 236–239
- geodesic triangle. *See* geodesic polygon
- Gergonne, Joseph Diaz, 115, 117, 147, 148, 152
- gingerbread man knot. *See* knot, gingerbread man
- Girard, Albert, 90–92, 227
- Goldbach, Christian, 1, 63, 66, 101
- golden ratio, 39
- granny knot. *See* knot, granny
- graph, x, xi, 5, 99–111, 119–129, 134–144, 154, 156, 157, 247, 284
- great circle, 87–89, 92, 93, 227
- great dodecahedron, 149, 150
- great icosahedron, 149–151
- great stellated dodecahedron, 57, 150, 151
- Gsell, Georg, 16
- Gsell, Katharina, 16, 24
- Gsell, Salome Abigail, 24
- Guthrie, Francis, 131–133
- Guthrie, Frederick, 132
- Hadamard, Jacques Salomon, 118
- Haeckel, Ernst, 33
- hairy ball theorem, 5, 209, 210, 214
- Haken, Wolfgang, 8, 142–144
- Hales, Thomas C., 143
- Halle University, 133
- Hamilton, Richard, 268, 269
- Hamilton, William Rowan, 132
- handle, 169, 170, 172, 179–181, 184, 246, 256
- Hardy, Godfrey Harold, 24, 123, 253
- Harriot, Thomas, 90–92, 227
- Harriot-Girard theorem, 90–96, 227
- Hauptvermutung*, 179, 262
- Heawood, Percy John, 138, 139

- hedra*, 27, 64, 66
Heegaard, Poul, 184
heliocentric astronomy. *See* Copernican astronomy
Hermocrates, 42
Hessel, Johann Friedrich Christian, 146–149, 174, 179, 245
Hierholzer, Carl, 105
Hilbert, David, 262, 268
Hippasus, 38–40
homeomorphism, 176, 179, 183, 184
homology, 254, 255, 258, 263, 264, 266
homology group, 263
Hopf, Heinz, 213, 214, 222
Hopf’s *Umlaufsatz*. *See* theorem of turning tangents
Hoppe, R., 154

icosahedron, 27, 32, 34, 36, 40, 42, 43, 46–48, 56, 58–61, 76–78, 260
index (of a zero of a vector field), 206–210, 212, 214, 215
indicatrix, 165
interior angle theorem, 221, 225–227
invariance of dimension theorem, 246
isotopic, 174, 186, 196
Ivan VI of Russia, 18
Ivanovna, Anna, 16, 18

Jamnitzer, Wentzel, 51, 54, 149
Jonquières, Ernest de, 84, 85, 145
Jordan, Marie Ennemond Camille, 118, 183
Jordan curve theorem, 166, 169, 231

Kant, Immanuel, 101
Kauffman, Louis, 199
Kelvin, Lord. *See* Thomson, William
Kempe, Alfred Bray, 137–142
Kempe chain, 139, 141, 142
Kepler, Johannes, x, 1, 8, 32, 49, 51–61, 92, 98, 143, 150, 260, 270; *Cosmic Mystery*, 55–57; *The Harmony of the World*, 57–59
Kepler conjecture, 143
Kepler-Poinsot polyhedron, 57, 58, 98, 149–151
Klein, Felix, 156, 159, 162, 165–169, 253, 262
Klein bottle, 167–169, 171, 178, 182, 209, 242, 245, 251, 257, 258, 273
knot, x, xii, 5, 186–201; figure eight, 186, 187, 193–197, 199, 200; gingerbread man, 186, 187, 193–200; granny, 196; pentafoil, 186, 187, 193–200; square, 186, 187, 191–200; trefoil, 185–200; unknot, 186, 187, 189, 190, 194–198, 200
knot invariant, 188, 189, 194, 197–200
Koch, John, 142
Koestler, Arthur, 53
Kroto, Harold, 7, 8
Kuhn, Thomas, 84, 86
Kuratowski’s reduction theorem, 124

Lagrange, Joseph-Louis, 21, 112, 243
Lakatos, Imre, 115, 117, 148, 284
Laplace, Pierre-Simon, 10, 112
Lebesgue, Henri, 72, 85, 141
Lefschetz, Solomon, 157–158, 213
Legendre, Adrien-Marie, 65, 67, 87–89, 92, 94–99, 145, 149, 150, 152, 156, 234; *Elements of Geometry*, 87
Leibniz, Gottfried, 14, 83, 91, 102, 103, 113
Leipzig University, 133
Leonard, K. C. von, 148
Leonardo da Vinci, 51, 53
Lhuillier, Simon-Antoine-Jean, 146–149, 152, 154, 170, 174, 179, 180
Listing, Johann Benedict, 108, 118, 157, 164, 188, 247–250, 253
Little, Charles Newton, 188
London, England, 14, 18. *See* Royal Society of London
London Mathematical Society, 132
lune, 92, 93

manifold, 214, 243–246, 252, 258–262, 265–268; with boundary, 244, 259, 262, 265
map coloring, 6, 7
Marinoni, Giovanni, 102
Marquis de Condorcet, 25
Mathematical Association of America, 142
May, Kenneth O., 131
Menelaus of Alexandria, 89; *Sphaerica*, 89
minimal criminal, 137, 138, 140–142

- Möbius, August, 133–135, 164, 165, 176, 183, 184, 247, 250, 253
- Möbius band, 3, 156, 162–165, 167, 169, 171, 176, 177, 180–182, 184, 188–190, 247, 250, 273
- Murasugi, Kunio, 195, 199
- music of the spheres, 38
- Nash, John Forbes Jr., 270
- Newton, Isaac, 10, 14, 16, 24, 57, 113
- Nobel Prize, 7, 12, 265
- Noether, Emmy Amalie, 262, 263
- nonorientable, 165–172, 177, 181, 183, 184, 190, 244, 246, 250, 256, 261
- normal vector, 166, 231, 232, 235
- octahedron, 32–34, 36, 40, 42, 46, 47, 56, 58–61, 75, 77, 78, 228, 229, 246, 247
- one-sided surface, 164–166, 190
- optics, 91
- orientable, 169, 170, 172, 177, 181–185, 190, 191, 193, 238, 239, 244, 245, 250, 251, 255, 256, 260, 261, 265
- Pacioli, Fra Luca, *De Divina Proportione*, 51, 54
- Pappus, 49, 50
- Paterson, Michael, 127
- pentafoil knot. *See* knot, pentafoil
- pentagram, 39, 150, 151
- pentakis dodecahedron, 53
- Perelman, Grigori, 9, 268–270
- perspective in artwork, 51
- Peter I of Russia, 11, 13–16, 18, 22, 23
- Peter II of Russia, 15, 16
- Peter III of Russia, 22
- Peter the Great. *See* Peter I of Russia
- phase space, 202–204
- Pick, Georg Alexander, 124, 125
- Pick's theorem, 124–126
- plane angle, 46, 47, 63, 84, 85, 225, 226
- Plato, 1, 8, 31, 32, 40–43, 51, 58; *Academy*, 41–42, 44; *Epinomis*, 31; *The Sophist*, 40; *Theaetetus*, 40; *Timaeus*, 42–43
- Platonic solid. *See* regular polyhedron
- Poincaré, Henri, x, 28, 75, 85, 155, 157, 158, 173, 211, 212, 241, 244, 245, 252–262, 264, 266, 270; *Analysis Situs*, 173, 253, 254, 258, 261
- Poincaré conjecture, 8, 9, 265–270, 285
- Poincaré duality, 260, 261
- Poincaré's dodecahedral space, 266
- Poincaré-Hopf theorem, 205, 208–211, 214–217
- Poincaré, Raymond, 211
- Poinsot, Louis, 97–99, 145, 149, 150, 152
- Pólya, George, 220
- Pont, Jean-Claude, 147
- prime knot, 195, 196, 198, 200
- Princeton University, 158, 213
- problem of the five princes, 133–135
- Proclus, 44, 45
- projective plane, 168, 169, 171, 172, 180, 182, 242, 251, 257, 258, 273
- pseudosphere, 236
- Ptolemaic astronomy, 55
- Ptolemy, 50
- pyramid, 27, 32, 34, 40, 67–72, 116, 152, 245, 259. *See* tetrahedron
- Pythagoras, x, 1, 31, 36–39, 45, 51, 270
- Pythagorean theorem, 8, 37, 125
- Pythagoreans, 36–40, 42, 150
- quadratic formula, 8
- quadratic reciprocity law, 234
- radiolaria, 33, 34
- Raleigh, Sir Walter, 91
- Raphael, *School of Athens*, 51, 52
- reducible configuration, 141, 142, 144
- regular polyhedron, xi, 8, 31–35, 39–43, 45–48, 51, 55–62, 75–78, 84, 86, 151, 174, 181, 227, 260, 273, 285; duality, 58–61, 260
- Riemann, Georg Friedrich Bernhard, x, 118, 181–183, 244, 248, 250–254, 258, 264, 270
- Riemann hypothesis, 267
- Riemann surface, 183
- Royal Society of London, 14, 21, 87, 139
- saddle (on a surface), 206, 227, 235–237
- saddle (zero of a vector field), 205–209
- San Marco Basilica, 52
- Schläfli, Ludwig, 168, 247, 259

- Seifert, Herbert, 190, 191
Seifert circle, 191
Seifert surface, 190–196
semiregular polyhedron, 49, 57, 174, 181
simple closed curve, 166, 169, 219, 222, 223, 231, 238
simple graph, 135, 136
simplex, 244, 245, 259, 260, 262
simplicial complex, 244, 245, 258, 259, 262
simply connected, 255, 256, 266, 268
sink (zero of a vector field), 205–209
six color theorem, 137, 140
Smale, Stephen, 267–269
small stellated dodecahedron, 57, 150, 151
Smalley, Richard, 7, 8
Socrates, 40–42, 51
Sommerville, D. M. Y., 145
source (zero of a vector field), 205, 206, 208, 209
sphere, 3, 4, 30, 38, 39, 46, 55–57, 61, 67, 80, 87–98, 118, 120, 143, 155–159, 166, 169–172, 174, 177–184, 192, 194, 204–206, 208–210, 212, 214, 217, 219, 225, 227, 236–239, 244, 246, 247, 249, 250, 254, 255, 258, 260; n -sphere (S^n), 212, 213, 217, 244, 247, 260, 266–268
sprouts, 127–129
square knot. *See* knot, square
star polyhedron. *See* Kepler-Poinsot polyhedron
star-convex polyhedron, 97–99, 145, 150
Staudt, Karl Georg Christian von, 152–155; *Geometrie der Lage*, 152
Steiner, Jakob, 152
Steinitz, Ernst, 117
stereoisomer, 48
surface, xi, xii, 5, 158–186, 189, 191, 204, 206, 208, 209, 212, 214–216, 219, 223, 227–230, 232, 233, 235–239, 241–246, 250, 251, 254–260, 265; with boundary, 159, 162, 167–169, 171, 177, 180, 181, 183–185, 189–196, 214, 217, 244, 250
Tait, Peter Guthrie, 157, 188, 189, 199, 249
tangent vector, 222, 223, 231
Terquem, Orly, 109
tetrahedron, 27, 31–34, 36, 40, 42, 46, 47, 56, 58–61, 64, 67, 68, 75, 77, 78, 224, 225
Thales, 31
Theaetetus, 1, 40–42, 44, 49, 86, 151, 181
Theodorus, 40
theorem of turning tangents, 222, 223, 232
Thistlethwaite, Morwen, 199
Thomson, William, 188, 189, 201; vortex model of atomic theory, 188, 201
Thurston, William, 214, 268, 269
Thurston’s geometrization conjecture, 268, 269, 285
topological invariant, 4, 177–179, 189, 225, 240, 241, 246, 250–252, 258–260
torsion coefficient, 257–261, 263, 266
torus, 3, 4, 148, 154, 156, 158–162, 167, 169, 170, 174–176, 178, 180, 182, 183, 185, 192, 193, 204–206, 208, 210, 225, 235, 239, 244, 245, 249–251, 255, 256, 258–260, 273; g -holed, 169, 170, 181, 182, 184, 192, 219, 258, 260; 3-torus, 245, 261; double, 161, 162, 169, 170, 174–176, 182, 183, 193, 208, 209, 251, 256, 258, 260, 266
tree, 120, 121, 153, 154
trefoil knot. *See* knot, trefoil
triangulate, 116, 184, 262
truncated icosahedron, 2, 27, 53
Tucker, Albert, 158
Uccello, Paolo, 52, 54, 149
unavoidable set, 141, 142, 144
University of Basel, 12, 15
University of Berlin, 181
University of Breslau, 213
University of Cambridge, 127
University of Erlangen, 152
University of Göttingen, 133, 152, 181, 234, 235, 251, 262
University of Pisa, 251
University of Würzburg, 152
unknot. *See* knot, unknot
Vandermonde, Alexandre-Théophile, 187
Veblen, Oswald, 141, 157
vector field, 203–210, 212, 214–217; gradient, 204–206, 208

Verona, Fra Giovanni da, 52, 53
Voltaire, 19, 20

Waterhouse, William, 34, 35, 40, 86
Weber, Wilhelm Eduard, 182
Whitney, Hassler, 141, 171, 245

Whitney embedding theorem, 246
Whitney umbrella, 171
Wiles, Andrew, 8, 270

Zeno, 31
zero of vector field, 204–210, 212, 214–218